

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Journal of Pure and Applied Algebra

journal homepage: www.elsevier.com/locate/jpaa

Hopf crossed module (co)algebras

Kürşat Sözer^a, Alexis Virelizier^{b,*}^a *McMaster University, Hamilton, ON, L8S 4E8, Canada*^b *Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé, Lille, F-59000, France*

ARTICLE INFO

Article history:

Received 29 May 2023

Received in revised form 18 March 2024

Available online 5 September 2025

Communicated by A. Solotar

MSC:

16T05; 18M05; 18A50; 18G45

Keywords:

Graded Hopf algebras

Crossed modules

Graded monoidal categories

Fusion categories

ABSTRACT

Given a crossed module χ , we introduce Hopf χ -(co)algebras which generalize Hopf algebras and Hopf group-(co)algebras. We interpret them as Hopf algebras in some symmetric monoidal category. We prove that their categories of representations are monoidal and χ -graded (meaning that both objects and morphisms have degrees which are related via χ).

© 2025 Elsevier B.V. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

1. Introduction

Hopf algebras, such as quantum groups, are fundamental objects in the field of quantum algebra and quantum topology. In particular, Hopf algebras and their categories of representations are very useful in the construction of quantum invariants of knots and 3-manifolds (see for instance [7,13,18,9,10,6,11]).

There are various generalizations of Hopf algebras. A particular generalization is given by so-called Hopf group-coalgebras introduced by Turaev for topological purposes: given a (discrete) group G , Hopf G -coalgebras and their categories of representations (which are G -graded monoidal categories) are used to define quantum invariants of principal G -bundles over 3-manifolds (see [16]). On the algebraic side, the second author generalized to Hopf group-coalgebras most of the classical results for Hopf algebras (see [19]), and Caenepeel and De Lombaerde showed that Hopf group-coalgebras are Hopf algebra objects in a certain symmetric monoidal category (see [4]).

In this paper, we introduce and study Hopf crossed module-coalgebras which are extensions of Hopf group-coalgebras. Recall from homotopical algebra that crossed modules are a convenient way of encoding

* Corresponding author.

E-mail addresses: sozerk@mcmaster.ca (K. Sözer), alexis.virelizier@univ-lille.fr (A. Virelizier).

(strict) 2-groups. Explicitly, a crossed module is a group homomorphism $\chi: E \rightarrow H$ together with an action of H on E such that χ is H -equivariant and satisfies the Peiffer identity (see Section 4.1).

Given a crossed module $\chi: E \rightarrow H$, a Hopf χ -coalgebra is a Hopf H -coalgebra endowed with a χ -action (see Section 7.2). More explicitly, a Hopf χ -coalgebra (over a commutative ring \mathbb{k}) is a family $A = \{A_x\}_{x \in H}$ of \mathbb{k} -algebras endowed with a comultiplication $\Delta = \{\Delta_{x,y}: A_{xy} \rightarrow A_x \otimes A_y\}_{x,y \in H}$, a counit $\varepsilon: A_1 \rightarrow \mathbb{k}$, an antipode $S = \{S_x: A_{x^{-1}} \rightarrow A_x\}_{x \in H}$, and a χ -action consisting of a family of algebra isomorphisms

$$\phi = \{\phi_{x,e}: A_x \rightarrow A_{\chi(e)x}\}_{(x,e) \in H \times E}.$$

These data should verify some compatibility conditions generalizing the axioms of a Hopf algebra and of an action of a group. There are two particular cases. First, if H is a group, then the trivial map $1 \rightarrow H$ is a crossed module and a Hopf $(1 \rightarrow H)$ -coalgebra is a Hopf H -coalgebra. Second, if E is an abelian group, then the map $E \rightarrow 1$ is a crossed module and a Hopf $(E \rightarrow 1)$ -coalgebra is a Hopf algebra endowed with an action of E by algebra and bicomodule automorphisms.

Our main motivation for introducing Hopf χ -coalgebras is to produce instances of monoidal categories which are χ -graded (in the sense of [14]). In such a category, not only the objects have a degree in H , but also the morphisms have a degree in E , and these two degrees are related by the crossed module homomorphism χ . Actually, χ -graded monoidal categories are useful for topological purposes: it is shown in [14] that any χ -graded χ -fusion category gives rise to a state sum invariant of 3-manifolds endowed with a homotopy class of maps to the classifying space $B\chi$ of χ (which is a homotopy 2-type) and, more generally, to a 3-dimensional homotopy quantum field theory with target $B\chi$.

The first main result of the paper is that the category $\text{Mod}_\chi(A)$ of modules over a Hopf χ -coalgebra A is a χ -graded monoidal category with internal Homs (see Theorem 8.1). We also study its full subcategory $\text{mod}_\chi(A)$ whose objects have their underlying module projective of finite rank. We prove that the pivotal structures on $\text{mod}_\chi(A)$ are in bijective correspondence with the pivotal elements of A (see Corollary 8.2), and we provide sufficient conditions on A for $\text{mod}_\chi(A)$ to be a χ -fusion category (see Theorem 8.3).

Note that the notion of a Hopf χ -coalgebra is not self-dual: the dual notion is that of a Hopf χ -algebra (see Section 7.8) and the category $\text{Comod}_\chi(A)$ of comodules over a Hopf χ -algebra A is a closed χ -graded monoidal category (see Section 8.8).

Next, we introduce the notion of a Hopf χ -module over a Hopf χ -coalgebra and prove a structure theorem for them (see Theorem 9.1). When the ground ring is a field and the Hopf χ -coalgebra is of finite type, we derive from this structure theorem the existence and uniqueness of χ -integrals (see Theorem 9.2). These generalize the well known corresponding results for Hopf algebras.

Finally, we interpret Hopf crossed module-(co)algebras in any symmetric monoidal category \mathcal{S} as Hopf algebra objects in some symmetric monoidal category associated with \mathcal{S} (see Theorems 10.1 and 10.2). In particular, the case where \mathcal{S} is the category $\text{Mod}_\mathbb{k}$ of \mathbb{k} -modules corresponds to the Hopf crossed module-(co)algebras over \mathbb{k} (considered above). This is built on the fact that crossed modules are group objects in the category of small categories (see [3]) and generalizes the above cited work of Caenepeel and De Lombaerde.

The paper is organized as follows. In Section 2, we review monoidal categories and the associated graphical calculus. In Section 3, we discuss the notions of categorical (co)algebras, Hopf algebras, and (co)modules. We recall crossed modules in Section 4 and crossed module graded categories in Section 5. In Section 6, we discuss Hopf group-coalgebras and their categories of representations. In Section 7, we introduce Hopf crossed module-(co)algebras. Section 8 is devoted to the study of their categories of representations. In Section 9, we introduce and characterize Hopf crossed module-modules and study integrals of Hopf crossed module-coalgebras. Finally, in Section 10, we interpret Hopf crossed module-(co)algebras as Hopf algebras in some symmetric monoidal category.

Throughout the paper, we fix a nonzero commutative ring \mathbb{k} . The tensor product over \mathbb{k} is denoted $\otimes_{\mathbb{k}}$ or more simply \otimes if there is no confusion.

2. Categorical preliminaries

2.1. Conventions on monoidal categories

For the basics on monoidal categories, we refer for example to [12,5,17]. Unless otherwise specified, we denote the monoidal product of a monoidal category by \otimes and its unit object by $\mathbb{1}$. We will suppress in our formulas the associativity and unitality constraints of monoidal categories. This does not lead to ambiguity because by Mac Lane’s coherence theorem, all legitimate ways of inserting these constraints give the same result. For any objects X_1, \dots, X_n with $n \geq 2$, we set

$$X_1 \otimes X_2 \otimes \cdots \otimes X_n = (\dots((X_1 \otimes X_2) \otimes X_3) \otimes \cdots \otimes X_{n-1}) \otimes X_n$$

and similarly for morphisms.

2.2. Braided and symmetric categories

A *braiding* on a monoidal category \mathcal{B} is a natural isomorphism $\tau = \{\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X\}_{X,Y \in \mathcal{B}}$ such that for all objects $X, Y, Z \in \mathcal{B}$,

$$\tau_{X,Y \otimes Z} = (\text{id}_Y \otimes \tau_{X,Z})(\tau_{X,Y} \otimes \text{id}_Z) \quad \text{and} \quad \tau_{X \otimes Y, Z} = (\tau_{X,Z} \otimes \text{id}_Y)(\text{id}_X \otimes \tau_{Y,Z}).$$

A *braided category* is a monoidal category endowed with a braiding.

A *braiding* on a monoidal category \mathcal{B} is *symmetric* if $\tau_{Y,X}\tau_{X,Y} = \text{id}_{X \otimes Y}$ for all $X, Y \in \mathcal{B}$. A *symmetry* is a symmetric braiding. A *symmetric monoidal category* is a monoidal category endowed with a symmetry.

2.3. Cartesian monoidal categories

A *cartesian monoidal category* is a monoidal category whose monoidal structure is given by the category-theoretic product (and so whose unit object is a terminal object). Such a category is then symmetric, with symmetry given by the canonical flip maps.

Any category with finite products can be considered as a cartesian monoidal category (as long as we have a specified product for each pair of objects). In particular, the category **Set** of sets and maps, and the category **Cat** of small categories and functors, endowed with their canonical category-theoretic product, are symmetric monoidal cartesian categories.

For any object X of a cartesian monoidal category, there is a unique morphism $\Delta_X : X \rightarrow X \otimes X$, called the *diagonal map*, such that Δ_X composed with the first or second projection is the identity and, since the unit object is a terminal object, there is a unique morphism $\varepsilon_X : X \rightarrow \mathbb{1}$, called the *augmentation*.

2.4. Rigid categories

A *duality* in a monoidal category \mathcal{C} is a quadruple (X, Y, e, d) , where X, Y are objects of \mathcal{C} , $e : X \otimes Y \rightarrow \mathbb{1}$ (the *evaluation*) and $d : \mathbb{1} \rightarrow Y \otimes X$ (the *coevaluation*) are morphisms in \mathcal{C} , such that

$$(e \otimes \text{id}_X)(\text{id}_X \otimes d) = \text{id}_X \quad \text{and} \quad (\text{id}_Y \otimes e)(d \otimes \text{id}_Y) = \text{id}_Y.$$

Then (X, e, d) is a *left dual* of Y and (Y, e, d) is a *right dual* of X .

Left and right duals, if they exist, are essentially unique: if (Y, e, d) and (Y', e', d') are right duals of some object X , then there exists a unique isomorphism $u: Y \rightarrow Y'$ such that $e' = e(\text{id}_X \otimes u^{-1})$ and $d' = (u \otimes \text{id}_X)d$.

A monoidal category is *left rigid* (respectively, *right rigid*) if every object admits a left dual (respectively, a right dual). A *rigid category* is a monoidal category which is both left rigid and right rigid.

Subsequently, when dealing with rigid categories, we shall always assume tacitly that for each object X , a left dual $({}^\vee X, \text{ev}_X, \text{coev}_X)$ and a right dual $(X^\vee, \tilde{\text{ev}}_X, \widetilde{\text{coev}}_X)$ has been chosen. Such a choice defines a left dual functor ${}^\vee?: \mathcal{C}^{\text{rev}} \rightarrow \mathcal{C}$ and a right dual functor $?^\vee: \mathcal{C}^{\text{rev}} \rightarrow \mathcal{C}$, where $\mathcal{C}^{\text{rev}} = (\mathcal{C}^{\text{op}}, \otimes^{\text{op}}, \mathbb{1})$ is the opposite category to \mathcal{C} with opposite monoidal structure. In particular, the left and right duals of any morphism $f: X \rightarrow Y$ in \mathcal{C} are defined by

$$\begin{aligned} {}^\vee f &= (\text{ev}_Y \otimes \text{id}_{{}^\vee X})(\text{id}_{{}^\vee Y} \otimes f \otimes \text{id}_{{}^\vee X})(\text{id}_{{}^\vee Y} \otimes \text{coev}_X): {}^\vee Y \rightarrow {}^\vee X, \\ f^\vee &= (\text{id}_{X^\vee} \otimes \tilde{\text{ev}}_Y)(\text{id}_{X^\vee} \otimes f \otimes \text{id}_{Y^\vee})(\widetilde{\text{coev}}_X \otimes \text{id}_{Y^\vee}): Y^\vee \rightarrow X^\vee. \end{aligned}$$

The left and right dual functors are monoidal. Note that the actual choice of left and right duals is innocuous in the sense that different choices of left (respectively, right) duals define canonically monoidally isomorphic left (respectively, right) dual functors.

2.5. Closed monoidal categories

A monoidal category \mathcal{C} is *left closed* if for any object X of \mathcal{C} , the endofunctor $? \otimes X$ has a right adjoint $[X, ?]^l$, with adjunction unit and counit:

$$\text{ev}_Y^X: [X, Y]^l \otimes X \rightarrow Y \quad \text{and} \quad \text{coev}_Y^X: Y \rightarrow [X, Y \otimes X]^l,$$

called respectively the *left evaluation* and the *left coevaluation*. Then $[X, Y]^l$ is called the *left internal Hom* from X to Y . The left internal Homs give rise to a functor $[-, -]^l: \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$.

Similarly, a monoidal category \mathcal{C} is *right closed* if for any object X of \mathcal{C} , the endofunctor $X \otimes ?$ has a right adjoint $[X, ?]^r$, with adjunction unit and counit:

$$\tilde{\text{ev}}_Y^X: X \otimes [X, Y]^r \rightarrow Y \quad \text{and} \quad \widetilde{\text{coev}}_Y^X: Y \rightarrow [X, X \otimes Y]^r,$$

called respectively the *right evaluation* and the *right coevaluation*. Then $[X, Y]^r$ is called the *right internal Hom* from X to Y . The right internal Homs give rise to a functor $[-, -]^r: \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$.

A monoidal category is *closed* if it is both left and right closed. For example, the category $\text{Mod}_{\mathbb{k}}$ of \mathbb{k} -modules and \mathbb{k} -linear homomorphisms is closed: the internal Homs between \mathbb{k} -modules M and N are $[M, N]^l = [M, N]^r = \text{Hom}_{\mathbb{k}}(M, N)$ with the standard (co)evaluations.

Any rigid monoidal category is closed: the internal Homs are $[X, Y]^l = Y \otimes {}^\vee X$ and $[X, Y]^r = X^\vee \otimes Y$ with (co)evaluations

$$\begin{aligned} \text{ev}_Y^X &= \text{id}_Y \otimes \text{ev}_X, & \text{coev}_Y^X &= \text{id}_Y \otimes \text{coev}_X, \\ \tilde{\text{ev}}_Y^X &= \tilde{\text{ev}}_X \otimes \text{id}_Y, & \widetilde{\text{coev}}_Y^X &= \widetilde{\text{coev}}_X \otimes \text{id}_Y. \end{aligned}$$

2.6. Pivotal categories

A *pivotal category* is a rigid category \mathcal{C} endowed with a monoidal isomorphism between the left and the right dual functors. By modifying the right duals of objects using this monoidal isomorphism, we may

assume it to be the identity without loss of generality. In other words, for each object X of \mathcal{C} , we have a dual object X^* and four morphisms

$$\begin{aligned} \text{ev}_X: X^* \otimes X &\rightarrow \mathbb{1}, & \text{coev}_X: \mathbb{1} &\rightarrow X \otimes X^*, \\ \widetilde{\text{ev}}_X: X \otimes X^* &\rightarrow \mathbb{1}, & \widetilde{\text{coev}}_X: \mathbb{1} &\rightarrow X^* \otimes X, \end{aligned}$$

such that $(X^*, \text{ev}_X, \text{coev}_X)$ is a left dual for X , $(X^*, \widetilde{\text{ev}}_X, \widetilde{\text{coev}}_X)$ is a right dual for X , and the induced left and right dual functors coincide as monoidal functors.

For example, any left rigid symmetric monoidal category has a canonical structure of a pivotal category, for which the right (co)evaluations are given by the left ones (pre)composed with the symmetry. In particular, the full subcategory of $\text{Mod}_{\mathbb{k}}$ consisting of projective \mathbb{k} -modules of finite rank has a canonical structure of a pivotal category: the dual of projective \mathbb{k} -module M of finite rank is $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$ with the standard (co)evaluations.

2.7. Penrose graphical calculus

Morphisms in a monoidal category may be represented by planar diagrams to be read from bottom to top. We discuss here the basics of this Penrose graphical calculus (see [8] or [17] for a detailed treatment). The diagrams are made up of arcs colored by objects and of boxes colored by morphisms. The arcs connect the boxes and have no intersections or self-intersections. The identity id_X of an object X , a morphism $f: X \rightarrow Y$, and the composition of two morphisms $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are represented as follows:

$$\text{id}_X = \left| \begin{array}{c} \\ \hline X \\ \hline \end{array} \right|, \quad f = \begin{array}{c} |Y \\ \boxed{f} \\ |X \end{array}, \quad \text{and} \quad gf = \begin{array}{c} |Z \\ \boxed{g} \\ |Y \\ \boxed{f} \\ |X \end{array}.$$

The monoidal product of two morphisms $f: X \rightarrow Y$ and $g: U \rightarrow V$ is represented by juxtaposition:

$$f \otimes g = \begin{array}{cc} |Y & |V \\ \boxed{f} & \boxed{g} \\ |X & |U \end{array}.$$


We can also use boxes with several strands attached to their horizontal sides. For example, a morphism $f: X \otimes Y \rightarrow A \otimes B \otimes C$ may be represented in various ways, such as

$$\begin{array}{c} |A \quad |B \quad |C \\ \boxed{f} \\ |X \quad |Y \end{array} \quad \text{or} \quad \begin{array}{c} |A \quad |B \otimes C \\ \boxed{f} \\ |X \otimes Y \end{array} \quad \text{or} \quad \begin{array}{c} A \otimes B \quad | \quad |C \\ \boxed{f} \\ |X \quad |Y \end{array}.$$

Here, in accordance with the conventions of Section 2.1, we ignore here the associativity constraint between the objects $A \otimes B \otimes C = (A \otimes B) \otimes C$ and $A \otimes (B \otimes C)$. A box whose lower/upper side has no attached strands represents a morphism with source/target $\mathbb{1}$. For example, morphisms $\alpha: \mathbb{1} \rightarrow \mathbb{1}$, $\beta: \mathbb{1} \rightarrow X$, $\gamma: X \rightarrow \mathbb{1}$ may be represented by the diagrams

$$\boxed{\alpha}, \quad \begin{array}{c} |X \\ \boxed{\beta} \end{array}, \quad \boxed{\gamma} \begin{array}{c} \\ |X \end{array}.$$

Every diagram which is colored as above determines a morphism obtained as follows. First slice the diagram into horizontal strips so that each strip is made of juxtaposition of vertical segments or boxes. Then, for each strip, take the monoidal product of the morphisms associated to the vertical segments or boxes. Finally, compose the resulting morphisms proceeding from the bottom to the top. For example, given morphisms $f: Y \rightarrow Z$, $g: B \otimes Z \rightarrow \mathbb{1}$, $h: X \rightarrow A \otimes B$, the diagram




represents the morphism

$$(\text{id}_A \otimes g \otimes \text{id}_X)(h \otimes f \otimes \text{id}_X) = ((\text{id}_A \otimes g)(h \otimes f)) \otimes \text{id}_X$$

from $X \otimes Y \otimes X$ to $A \otimes X$.

The functoriality of the monoidal product implies that the morphism associated to a colored diagram is independent of the way of cutting it into horizontal strips. It also implies that we can push boxes lying on the same horizontal level up or down without changing the morphism represented by the diagram. For example, for all morphisms $f: X \rightarrow Y$ and $g: U \rightarrow V$, we have:



which graphically expresses the formulas

$$f \otimes g = (\text{id}_Y \otimes g)(f \otimes \text{id}_U) = (f \otimes \text{id}_V)(\text{id}_X \otimes g).$$

Here and in the sequel, the equality sign between the diagrams means the equality of the corresponding morphisms.


The braiding τ of a braided category, and its inverse, are depicted as

$$\tau_{X,Y} = \begin{array}{c} Y \quad X \\ \diagdown \quad / \\ X \quad Y \end{array} \quad \text{and} \quad \tau_{Y,X}^{-1} = \begin{array}{c} Y \quad X \\ / \quad \diagdown \\ X \quad Y \end{array}.$$

The axioms of a braiding are depicted as follows: for all objects X, Y, Z ,

$$\begin{array}{c} Z \quad X \otimes Y \\ \diagdown \quad / \\ X \otimes Y \quad Z \end{array} = \begin{array}{c} Z \quad X \quad Y \\ \diagdown \quad / \quad \diagdown \quad / \\ X \quad Y \quad Z \quad Z \end{array} \quad \text{and} \quad \begin{array}{c} Y \otimes Z \quad X \\ \diagdown \quad / \\ X \quad Y \otimes Z \end{array} = \begin{array}{c} Y \quad Z \quad X \\ \diagdown \quad / \quad \diagdown \quad / \\ X \quad Y \quad Z \quad Z \end{array}.$$


The naturality of τ ensures that we can push boxes across a strand without changing the morphism represented by the diagram: for any morphism f ,



The above graphical calculus may be enhanced for a pivotal category by orienting all arcs in the diagrams and depicting the (co)evaluations as

$$\text{ev}_X = \text{arc}_X^{\downarrow}, \quad \tilde{\text{ev}}_X = \text{arc}_X^{\uparrow}, \quad \text{coev}_X = \text{arc}_X^{\downarrow}, \quad \widetilde{\text{coev}}_X = \text{arc}_X^{\uparrow}.$$


Here, an arc colored with an object X and oriented downward (resp., upward) contributes X (resp., X^*) to the source/target of morphisms. For example, a morphism $f: X^* \otimes Y \rightarrow A \otimes B^* \otimes C$ may be represented as



In particular, the identity of the dual X^* of any object X is represented as

$$\text{id}_{X^*} = \text{downward arc}_X = \text{upward arc}_X.$$

The duality identities are graphically expressed as



In a pivotal category, the morphism represented by a diagram is preserved under ambient isotopies of the diagram keeping fixed the bottom and top endpoints.

3. Categorical Hopf algebras


In this section, we review the categorical version of the notions of a Hopf algebra and its modules.

3.1. Categorical algebras


Let \mathcal{C} be a monoidal category. An *algebra in \mathcal{C}* is an object A of \mathcal{C} endowed with morphisms $m: A \otimes A \rightarrow A$ (the product) and $u: \mathbb{1} \rightarrow A$ (the unit) such that

$$m(m \otimes \text{id}_A) = m(\text{id}_A \otimes m) \quad \text{and} \quad m(\text{id}_A \otimes u) = \text{id}_A = m(u \otimes \text{id}_A).$$

We depict the product and the unit as



The axioms above have the following graphical interpretation:



Note that algebras in \mathcal{C} are also called *monoids* in \mathcal{C} in the literature.

3.2. Categorical coalgebras

A *coalgebra* in a monoidal category \mathcal{C} is an algebra in the opposite category $\mathcal{C}^{\text{op}} = (\mathcal{C}^{\text{op}}, \otimes, \mathbb{1})$. In other words, a coalgebra in \mathcal{C} is an object A of \mathcal{C} endowed with morphisms $\Delta: A \rightarrow A \otimes A$ (the coproduct) and $\varepsilon: A \rightarrow \mathbb{1}$ (the counit) such that

$$(\Delta \otimes \text{id}_A)\Delta = (\text{id}_A \otimes \Delta)\Delta \quad \text{and} \quad (\text{id}_A \otimes \varepsilon)\Delta = \text{id}_A = (\varepsilon \otimes \text{id}_A)\Delta.$$

We depict the coproduct and the counit as

$$\Delta = \begin{array}{c} A \quad A \\ \cup \\ A \end{array} \quad \text{and} \quad \varepsilon = \begin{array}{c} \circ \\ | \\ A \end{array}.$$

The axioms above are depicted as

$$\begin{array}{c} A \quad A \quad A \\ \cup \quad \cup \\ A \end{array} = \begin{array}{c} A \quad A \quad A \\ \cup \quad \cup \\ A \end{array} \quad \text{and} \quad \begin{array}{c} A \\ \cup \quad \circ \\ A \end{array} = \begin{array}{c} A \\ | \\ A \end{array} = \begin{array}{c} \circ \\ | \\ A \end{array}.$$

For example, in a cartesian monoidal category (see Section 2.3), any object is a coalgebra with coproduct being the diagonal map and counit being the augmentation. In fact, any coalgebra in a cartesian monoidal category is of this form.

3.3. Categorical bialgebras

To define bialgebras in a monoidal category, we need compatibility conditions between product and coproduct, and the formulation of one of the conditions requires a substitute for the flip map which can be provided by a braiding. A *bialgebra* in a braided category \mathcal{B} is an object A of \mathcal{B} endowed with an algebra structure (m, u) and a coalgebra structure (Δ, ε) in \mathcal{B} satisfying the following conditions (expressing that Δ and ε are algebra morphisms or, equivalently, that m and u are coalgebra morphisms):

$$\begin{aligned} \Delta m &= (m \otimes m)(\text{id}_A \otimes \tau_{A,A} \otimes \text{id}_A)(\Delta \otimes \Delta), & \Delta u &= u \otimes u, \\ \varepsilon m &= \varepsilon \otimes \varepsilon, & \varepsilon u &= \text{id}_{\mathbb{1}}, \end{aligned}$$

where τ is the braiding of \mathcal{B} . Pictorially,


$$\begin{array}{c} A \quad A \\ \cup \\ A \quad A \end{array} = \begin{array}{c} A \quad A \\ \cup \quad \cup \\ A \quad A \end{array}, \quad \begin{array}{c} A \quad A \\ \cup \\ \circ \end{array} = \begin{array}{c} A \quad A \\ | \quad | \\ \circ \quad \circ \end{array}, \quad \begin{array}{c} \circ \\ \cup \\ A \quad A \end{array} = \begin{array}{c} \circ \quad \circ \\ | \quad | \\ A \quad A \end{array}, \quad \begin{array}{c} \circ \\ | \\ A \end{array} = \emptyset.$$

3.4. Categorical Hopf algebras

Let \mathcal{B} be a braided category. An *antipode* for a bialgebra $A = (A, m, u, \Delta, \varepsilon)$ in \mathcal{B} is a morphism $S: A \rightarrow A$ such that

$$m(S \otimes \text{id}_A)\Delta = u\varepsilon = m(\text{id}_A \otimes S)\Delta.$$


This axiom is depicted as




If it exists, an antipode is unique and is anti-multiplicative and anti-comultiplicative:

$$Sm = m(S \otimes S)\tau_{A,A}, \quad Su = u, \quad \Delta S = \tau_{A,A}(S \otimes S)\Delta, \quad \varepsilon S = \varepsilon.$$

Pictorially,



When the antipode S is invertible, we depict its inverse $S^{-1}: A \rightarrow A$ as



A Hopf algebra in \mathcal{B} is a bialgebra in \mathcal{B} which admits an invertible antipode. Note that the notion of a Hopf algebra is self-dual: $(A, m, u, \Delta, \varepsilon, S)$ is a Hopf algebra in \mathcal{B} if and only if $(A, \Delta, \varepsilon, m, u, S)$ is a Hopf algebra in the opposite category \mathcal{B}^{op} .

A grouplike element of Hopf algebra A in \mathcal{B} is a morphism $G: \mathbb{1} \rightarrow A$ such that

$$\Delta G = G \otimes G \quad \text{and} \quad \varepsilon G = \text{id}_{\mathbb{1}}.$$

Such a G is invertible in the monoid $(\text{Hom}_{\mathcal{B}}(\mathbb{1}, A), *, u)$, where $\alpha * \beta = m(\alpha \otimes \beta)$, and its inverse, denoted by G^{-1} , is also a grouplike element of A and is computed by $G^{-1} = SG = S^{-1}G$. In particular, the set of grouplike elements of A is a group (with product $*$ and unit u).

3.5. Examples

1. Hopf algebras in the symmetric monoidal category of \mathbb{k} -modules and \mathbb{k} -linear homomorphisms are the usual Hopf \mathbb{k} -algebras.

2. Hopf algebras in the symmetric monoidal category of super \mathbb{k} -modules and grading-preserving \mathbb{k} -linear homomorphisms are the usual super Hopf \mathbb{k} -algebras.


3. Any group object in a cartesian monoidal category becomes a Hopf algebra, with its canonical coalgebra structure (see Section 3.2) and with antipode given by the group inversion. This induces a bijective correspondence between group objects and Hopf algebras in a cartesian monoidal category. For example, Hopf algebras in **Set** are groups. More generally, Hopf algebras in **Cat** are crossed modules, as detailed in Section 4.3.

3.6. Modules in categories

Let (A, m, u) be an algebra in a monoidal category \mathcal{C} . A *left A -module* (in \mathcal{C}) is a pair (M, r) , where M is an object of \mathcal{C} and $r: A \otimes M \rightarrow M$ is a morphism in \mathcal{C} , such that


$$r(m \otimes \text{id}_M) = r(\text{id}_A \otimes r) \quad \text{and} \quad r(u \otimes \text{id}_M) = \text{id}_M.$$

Graphically, these conditions are depicted as



One can similarly introduce right A -modules, but we will not use them. From now on, by an A -module, we mean a left A -module.

An *A -linear morphism* between two A -modules (M, r) and (N, s) is a morphism $f: M \rightarrow N$ in \mathcal{C} such that $fr = s(\text{id}_A \otimes f)$, that is, pictorially,



We let $\text{Mod}_{\mathcal{C}}(A)$ be the category of A -modules and A -linear morphisms, with composition inherited from \mathcal{C} . The *forgetful functor* $\text{Mod}_{\mathcal{C}}(A) \rightarrow \mathcal{C}$ carries any A -module (M, r) to M and any A -linear morphism to itself.

If $A = (A, m, u, \Delta, \varepsilon)$ is a bialgebra in a braided category \mathcal{B} , then the category $\text{Mod}_{\mathcal{B}}(A)$ of A -modules has a canonical structure of a monoidal category. Its unit object is the pair $(\mathbb{1}, \varepsilon)$. Its monoidal product is given on the objects by

$$(M, r) \otimes (N, s) = (M \otimes N, t)$$

where

$$t = (r \otimes s)(\text{id}_A \otimes \tau_{A, M} \otimes \text{id}_N)(\Delta \otimes \text{id}_{M \otimes N}) = \text{diagram}$$

and on the morphisms by the monoidal product in \mathcal{C} . Note that the forgetful functor $\text{Mod}_{\mathcal{B}}(A) \rightarrow \mathcal{B}$ is strict monoidal.

Assume that \mathcal{B} is a closed braided category. Then a bialgebra A in \mathcal{B} is a Hopf algebra if and only if the monoidal category $\text{Mod}_{\mathcal{B}}(A)$ is closed and the forgetful functor $\text{Mod}_{\mathcal{B}}(A) \rightarrow \mathcal{B}$ preserves the internal Homs. (This follows from Theorem 3.6 and Remark 5.6 of [1].) If such is the case, then the left and right internal Homs between A -modules (M, r) and (N, s) are

$$\begin{aligned} [(M, r), (N, s)]^l &= \left([M, N]^l, [\text{id}_M, \Theta_l]^l \text{coev}_{A \otimes [M, N]^l}^M \right), \\ [(M, r), (N, s)]^r &= \left([M, N]^r, [\text{id}_M, \Theta_r]^r \widetilde{\text{coev}}_{A \otimes [M, N]^r}^M \right), \end{aligned}$$

$$(M, r)^* = (M^*, r^\dagger) \quad \text{where} \quad r^\dagger = \begin{array}{c} \text{---} M \text{---} \\ \oplus \\ \text{---} A \text{---} \\ \uparrow M \end{array},$$

with left (co)evaluations given by $\text{ev}_{(M,r)} = \text{ev}_M$ and $\text{coev}_{(M,r)} = \text{coev}_M$, and right (co)evaluations given by

$$\widetilde{\text{ev}}_{(M,r)} = \begin{array}{c} A \\ \curvearrowright \\ \boxed{G} \\ \downarrow \\ \boxed{\gamma_M} \\ \downarrow M \end{array} \quad \text{and} \quad \widetilde{\text{coev}}_{(M,r)} = \begin{array}{c} \uparrow M \\ \boxed{\gamma_M^{-1}} \\ \downarrow \\ \boxed{G^{-1}} \\ \curvearrowleft \\ A \end{array}.$$

Note that the forgetful functor $\text{Mod}_{\mathcal{B}}(A) \rightarrow \mathcal{B}$ is then pivotal if and only if for all A -module (M, r) ,

$$\begin{array}{c} \uparrow M \\ \boxed{G} \\ \downarrow \\ \boxed{\gamma_M} \\ \downarrow M \end{array} = \begin{array}{c} \downarrow M \end{array}.$$

In particular, if A is *involutory* in the sense that its antipode S satisfies $S^2 = \theta_A$, then the monoidal category $\text{Mod}_{\mathcal{B}}(A)$ carries a structure of a pivotal category so that the forgetful functor $\text{Mod}_{\mathcal{B}}(A) \rightarrow \mathcal{B}$ is pivotal (by taking $G = u$ and $\gamma = \text{id}_{\mathcal{B}}$).

3.7. Comodules in categories

Given a coalgebra A in a monoidal category \mathcal{C} (which is an algebra in the opposite category \mathcal{C}^{op}), we define the category of left A -comodules in \mathcal{C} by setting

$$\text{Comod}_{\mathcal{C}}(A) = (\text{Mod}_{\mathcal{C}^{\text{op}}}(A))^{\text{op}}.$$

In particular its objects are pairs (M, δ) , with M an object of \mathcal{C} and $\delta: M \rightarrow A \otimes M$ a morphism in \mathcal{C} such that

$$(\Delta \otimes \text{id}_M)\delta = (\text{id}_A \otimes \delta)\delta \quad \text{and} \quad (\varepsilon \otimes \text{id}_M)\delta = \text{id}_M.$$

We deduce from Section 3.6 that the category $\text{Comod}_{\mathcal{C}}(A)$ is monoidal when \mathcal{C} is braided and A is a bialgebra, and that $\text{Comod}_{\mathcal{C}}(A)$ is rigid (resp. closed) when \mathcal{C} is braided rigid (resp. braided closed) and A is a Hopf algebra. Also, if \mathcal{C} is pivotal braided and A is a Hopf algebra, then the pivotal structures on the monoidal category $\text{Comod}_{\mathcal{C}}(A)$ are in bijection with the pairs (α, γ) , where $\alpha: A \rightarrow \mathbb{1}$ is an algebra morphism and $\gamma = \{\gamma_X: X \rightarrow X\}_{X \in \mathcal{B}}$ is a monoidal natural automorphism, such that $S^2 = \theta_A \circ \text{Ad}_{\alpha} \circ \gamma_A$, where θ is the twist of \mathcal{C} and

$$\text{Ad}_{\alpha} = \begin{array}{c} A \\ \downarrow \\ \boxed{\alpha} \\ \downarrow \\ \oplus \\ \downarrow \\ \boxed{\alpha} \\ \downarrow \\ A \end{array}.$$

4. Crossed modules

In this section, we quickly review crossed modules and their relationship with Hopf algebras in the category of small categories.

4.1. Crossed modules

A *crossed module* is a group homomorphism $\chi: E \rightarrow H$ together with a left action of H on E (by group automorphisms) denoted

$$(x, e) \in H \times E \mapsto {}^x e \in E$$

such that χ is equivariant with respect to the conjugation action of H on itself and satisfies the Peiffer identity, that is, for all $x \in H$ and $e, f \in E$,

$$\chi({}^x e) = x\chi(e)x^{-1} \quad \text{and} \quad \chi({}^e f) = e f e^{-1}.$$

These axioms imply that the image $\text{Im}(\chi)$ is normal in H and that the kernel $\text{Ker}(\chi)$ is central in E and is acted on trivially by $\text{Im}(\chi)$. In particular, $\text{Ker}(\chi)$ inherits an action of $H/\text{Im}(\chi) = \text{Coker}(\chi)$.

A *morphism* from a crossed module $\chi: E \rightarrow H$ to another crossed module $\chi': E' \rightarrow H'$ is a pair $(\psi: E \rightarrow E', \varphi: H \rightarrow H')$ of group homomorphisms such that

$$\chi'(\psi(e)) = \varphi(\chi(e)) \quad \text{and} \quad \psi({}^x e) = {}^{\varphi(x)}\psi(e)$$

for all $e \in E$ and $x \in H$.

4.2. Examples

1. Given any normal subgroup E of a group H , the inclusion $E \hookrightarrow H$ is a crossed module with the conjugation action of H on E .
2. For any group E , the homomorphism $E \rightarrow \text{Aut}(E)$ sending any element of E to the corresponding inner automorphism is a crossed module.
3. If E is an abelian group, then the trivial map $E \rightarrow 1$ is a crossed module.
4. A key geometric example of a crossed module is due to Whitehead: if X is a topological space, Y is a subspace of X , and y is a point of Y , then the homotopy boundary map $\partial: \pi_2(X, Y, y) \rightarrow \pi_1(Y, y)$, together with the standard action of $\pi_1(Y, y)$ on $\pi_2(X, Y, y)$, is a crossed module.

4.3. Crossed modules as Hopf algebras

By [3], crossed modules are group objects in the category **Cat** of small categories and functors (endowed with its canonical cartesian monoidal structure). Also, by the third example in Section 3.5, there is a bijective correspondence between group objects and Hopf algebras in a cartesian monoidal category. Consequently, Hopf algebras in **Cat** are in bijective correspondence with crossed modules.

Explicitly, the Hopf algebra $\mathcal{G}_\chi = (\mathcal{G}_\chi, m_\chi, u_\chi, \Delta_\chi, \varepsilon_\chi, S_\chi)$ in **Cat** associated with a crossed module $\chi: E \rightarrow H$ is described as follows. The objects of \mathcal{G}_χ are the elements of H . For any objects $x, y \in H$,

$$\text{Hom}_{\mathcal{G}_\chi}(x, y) = \{e \in E \mid y = \chi(e)x\}.$$

The composition of morphisms is given by the product of E :

$$(y \xrightarrow{f} z) \circ (x \xrightarrow{e} y) = (x \xrightarrow{fe} z) \quad \text{and} \quad \text{id}_x = 1 \in E.$$

Note that \mathcal{G}_χ is a groupoid. The product $m_\chi: \mathcal{G}_\chi \times \mathcal{G}_\chi \rightarrow \mathcal{G}_\chi$ of \mathcal{G}_χ is defined on objects and morphisms by

$$m_\chi(x, y) = xy \quad \text{and} \quad m_\chi(x \xrightarrow{e} y, z \xrightarrow{f} t) = (xz \xrightarrow{e^x f} yt).$$

Denote by $\mathbf{1}$ the trivial category with a single object $*$ and a single morphism id_* . The unit $u_\chi: \mathbf{1} \rightarrow \mathcal{G}_\chi$ is defined by

$$u_\chi(*) = 1 \in H.$$

The coproduct $\Delta_\chi: \mathcal{G}_\chi \rightarrow \mathcal{G}_\chi \times \mathcal{G}_\chi$ and counit $\varepsilon_\chi: \mathcal{G}_\chi \rightarrow \mathbf{1}$ are the diagonal and augmentation: for any object x and morphism e ,

$$\Delta_\chi(x) = (x, x), \quad \Delta_\chi(e) = (e, e), \quad \varepsilon_\chi(x) = *, \quad \varepsilon_\chi(e) = \text{id}_*.$$

The antipode $S_\chi: \mathcal{G}_\chi \rightarrow \mathcal{G}_\chi$ is involutive ($S_\chi^{-1} = S_\chi$) and is computed by

$$S_\chi(x) = x^{-1} \quad \text{and} \quad S_\chi(x \xrightarrow{e} y) = \left(x^{-1} \xrightarrow{x^{-1}(e^{-1})} y^{-1} \right).$$

5. Graded monoidal categories

In this section, we review the notions of a monoidal category graded by a group or a crossed module. We refer to [14] for details.

5.1. Linear categories

A category \mathcal{C} is \mathbb{k} -linear if for all objects $X, Y \in \mathcal{C}$, the set $\text{Hom}_{\mathcal{C}}(X, Y)$ carries a structure of a left \mathbb{k} -module so that the composition of morphisms is \mathbb{k} -bilinear. An object X of a \mathbb{k} -linear category \mathcal{C} is called a zero object if $\text{id}_X = 0$. A zero object, if it exists, is unique up to isomorphism.

A monoidal category is \mathbb{k} -linear if it is \mathbb{k} -linear as a category and the monoidal product of morphisms is \mathbb{k} -bilinear.

5.2. Monoidal categories graded by a group

Let H be a group. An H -graded monoidal category (over \mathbb{k}) is a \mathbb{k} -linear monoidal category \mathcal{C} endowed with a family $\{\mathcal{C}_h\}_{h \in H}$ of full subcategories such that:

- (a) Each object of \mathcal{C} is a direct sum of objects in $\bigcup_{h \in H} \mathcal{C}_h$.
- (b) For all $X \in \mathcal{C}_h$ and $Y \in \mathcal{C}_k$ with $h \neq k$,

$$\text{Hom}_{\mathcal{C}}(X, Y) = 0.$$

- (c) For all $X \in \mathcal{C}_h$ and $Y \in \mathcal{C}_k$, we have: $X \otimes Y \in \mathcal{C}_{hk}$.
- (d) $\mathbb{1} \in \mathcal{C}_1$.

In this case, we write by abuse¹ of notation:

$$\mathcal{C} = \bigoplus_{h \in H} \mathcal{C}_h.$$

The monoidal subcategory \mathcal{C}_1 is called the *neutral component* of \mathcal{C} . An object X of \mathcal{C} is *homogeneous* if it is nonzero and $X \in \mathcal{C}_h$ for some $h \in H$. Such an h is then uniquely determined by X , denoted by $|X|$, and called the *degree* of X .

5.3. Hom-graded categories

Let E be a group with unit 1. An E -Hom-graded category (over \mathbb{k}) is a category enriched over the monoidal category of E -graded \mathbb{k} -modules and \mathbb{k} -linear grading-preserving homomorphisms. Explicitly, this is a \mathbb{k} -linear category \mathcal{C} such that:

(a) The Hom-sets in \mathcal{C} are E -graded \mathbb{k} -modules: for all objects $X, Y \in \mathcal{C}$,

$$\text{Hom}_{\mathcal{C}}(X, Y) = \bigoplus_{e \in E} \text{Hom}_{\mathcal{C}}^e(X, Y).$$

(b) The composition in \mathcal{C} is multiplicative with respect to the degree: for all $e, f \in E$ and $X, Y, Z \in \mathcal{C}$, it sends $\text{Hom}_{\mathcal{C}}^f(Y, Z) \times \text{Hom}_{\mathcal{C}}^e(X, Y)$ into $\text{Hom}_{\mathcal{C}}^{fe}(X, Z)$.

(c) The identities have trivial degree: for all $X \in \mathcal{C}$,

$$\text{id}_X \in \text{End}_{\mathcal{C}}^1(X) = \text{Hom}_{\mathcal{C}}^1(X, X).$$

Let \mathcal{C} be an E -Hom-graded category. A morphism $\alpha: X \rightarrow Y$ in \mathcal{C} is *homogeneous of degree* $e \in E$ if $\alpha \in \text{Hom}_{\mathcal{C}}^e(X, Y)$. Note that if α is nonzero, then such an $e \in E$ is unique, is called the *degree* of α , and is denoted $e = |\alpha|$. The objects of \mathcal{C} together with the homogeneous morphisms of degree 1 form a \mathbb{k} -linear subcategory of \mathcal{C} called the *1-subcategory* of \mathcal{C} and denoted \mathcal{C}^1 .

Given $e \in E$, by an e -isomorphism we mean an isomorphism which is homogeneous of degree e . We say that an object X is e -isomorphic to an object Y if there is an e -isomorphism $X \rightarrow Y$.

Given $e \in E$, an object D of \mathcal{C} is an e -direct sum of a finite family $(X_a)_{a \in A}$ of objects of \mathcal{C} if there is a family $(p_a, q_a)_{a \in A}$ of morphisms such that for all $a, b \in A$, $p_a: D \rightarrow X_a$ is homogeneous of degree e^{-1} , $q_a: X_a \rightarrow D$ is homogeneous of degree e , $p_a q_b = \delta_{a,b} \text{id}_{X_a}$, and $\text{id}_D = \sum_{a \in A} q_a p_a$. Such an e -direct sum D , if it exists, is unique up to a 1-isomorphism and is denoted by

$$D = \bigoplus_{a \in A}^e X_a.$$

Note that for any finite families $(X_a)_{a \in A}$ and $(Y_b)_{b \in B}$ of objects of \mathcal{C} and for any $d, e, f \in E$, there are \mathbb{k} -linear isomorphisms

$$\text{Hom}_{\mathcal{C}}^d \left(\bigoplus_{a \in A}^e X_a, \bigoplus_{b \in B}^f Y_b \right) \cong \bigoplus_{\substack{a \in A \\ b \in B}} \text{Hom}_{\mathcal{C}}^{f^{-1}de} (X_a, Y_b). \tag{1}$$

¹ This is a genuine direct sum when \mathcal{C} is additive and each \mathcal{C}_h contains a zero object.

By definition, a direct sum of an empty family of objects of a \mathbb{k} -linear category \mathcal{C} is a *zero object* of \mathcal{C} , that is, an object $\mathbf{0}$ of \mathcal{C} such that $\text{End}_{\mathcal{C}}(\mathbf{0}) = 0$.

An E -Hom-graded category \mathcal{C} is *E-additive* if any finite (possibly empty) family of objects of \mathcal{C} has an e -direct sum in \mathcal{C} for all $e \in E$.

5.4. *Monoidal categories graded by a crossed module*

Let $\chi: E \rightarrow H$ be a crossed module. A χ -graded monoidal category (over \mathbb{k}), or shorter a χ -category, is a \mathbb{k} -linear monoidal category \mathcal{C} such that:

- (a) The \mathbb{k} -linear category \mathcal{C} is E -Hom-graded (see Section 5.3).
- (b) The associativity constraints $(X \otimes Y) \otimes Z \cong X \otimes (Y \otimes Z)$ and the unitality constraints $X \otimes \mathbf{1} \cong X \cong \mathbf{1} \otimes X$ of \mathcal{C} are all homogeneous of degree $1 \in E$.
- (c) The 1-subcategory \mathcal{C}^1 of \mathcal{C} (endowed with the monoidal structure induced by \mathcal{C}) is H -graded (see Section 5.2).

These data should satisfy two conditions relating the degree of objects and morphisms. To express them, we say that an object X of \mathcal{C} is *homogeneous* if it is homogeneous in \mathcal{C}^1 (see Section 5.2). Recall that the degree of a homogeneous object X is denoted by $|X| \in H$ and that the degree of a nonzero homogeneous morphism α is denoted by $|\alpha| \in E$. The two conditions are:

- (d) For all homogeneous objects X, Y and $e \in E$ such that $|Y| \neq \chi(e)|X|$,

$$\text{Hom}_{\mathcal{C}}^e(X, Y) = 0.$$

- (e) The monoidal product $\alpha \otimes \beta$ of any two nonzero homogeneous morphisms α, β is a homogeneous morphism of degree

$$|\alpha \otimes \beta| = |\alpha|^{s(\alpha)}|\beta|,$$

whenever the source $s(\alpha)$ of α is a homogeneous object. In other words, for any objects X, Y, Z, T with X homogeneous and for any morphisms $\alpha \in \text{Hom}_{\mathcal{C}}^e(X, Y)$, $\beta \in \text{Hom}_{\mathcal{C}}^f(Z, T)$ with $e, f \in E$, we have:

$$\alpha \otimes \beta \in \text{Hom}_{\mathcal{C}}^{e|X|f}(X \otimes Z, Y \otimes T).$$

Observe that this definition of a χ -category is equivalent to the definition given in [14] (by taking \mathcal{C}_{hom} to be the class of homogeneous objects of \mathcal{C} and the degree map $|\cdot|: \mathcal{C}_{\text{hom}} \rightarrow H$ to be the degree of homogeneous objects).

Note that the convention of Section 2.1 remains valid since Axiom (b) implies that the suppression of the associativity and unitality constraints does not change the degree of morphisms. Also, it follows from the axioms of an H -graded category that each object of \mathcal{C} is a 1-direct sum of a finite family of homogeneous objects. In particular, the Hom-sets in \mathcal{C} are fully determined by the Hom-sets between homogeneous objects. Axiom (d) implies that for any homogeneous objects X, Y ,

$$\text{Hom}_{\mathcal{C}}(X, Y) = \bigoplus_{e \in \chi^{-1}(|Y||X|^{-1})} \text{Hom}_{\mathcal{C}}^e(X, Y) \quad \text{and} \quad \text{End}_{\mathcal{C}}(X) = \bigoplus_{e \in \text{Ker}(\chi)} \text{End}_{\mathcal{C}}^e(X).$$

In particular $\text{Hom}_{\mathcal{C}}(X, Y) = 0$ whenever $|Y||X|^{-1} \notin \text{Im}(\chi)$. Also, if $\alpha: X \rightarrow Y$ is a nonzero homogeneous morphism between homogeneous objects, then

$$|Y| = \chi(|\alpha|) |X|.$$

In particular, 1-isomorphic homogeneous objects have the same degree.

A χ -category \mathcal{C} is *closed* if it is closed as a monoidal category (see Section 2.5) and all (co)evaluations are homogeneous of degree $1 \in E$. Note that a χ -category \mathcal{C} is closed if and only if its 1-subcategory \mathcal{C}^1 is closed.

A χ -category \mathcal{C} is *rigid* if it is rigid as a monoidal category (see Section 2.4) and all (co)evaluations are homogeneous of degree $1 \in E$. Note that a χ -category \mathcal{C} is rigid if and only if its 1-subcategory \mathcal{C}^1 is rigid.

A χ -category \mathcal{C} is *pivotal* if it is endowed with a pivotal structure (see Section 2.6) such that the dual X^* of any homogeneous object X is a homogeneous object of degree $|X^*| = |X|^{-1}$ and all (co)evaluations are homogeneous of degree $1 \in E$. Note that a χ -category \mathcal{C} is pivotal if and only if its 1-subcategory \mathcal{C}^1 is pivotal.

5.5. Example

The linearization $\mathbb{k}\mathcal{G}_\chi$ of the groupoid \mathcal{G}_χ associated with a crossed module $\chi: E \rightarrow H$ (see Section 4.3) is a χ -graded monoidal category. The objects of $\mathbb{k}\mathcal{G}_\chi$ are the elements of H . Each $x \in H$ is homogeneous with degree $|x| = x$. For any $x, y \in H$ and $e \in E$,

$$\text{Hom}_{\mathbb{k}\mathcal{G}_\chi}^e(x, y) = \begin{cases} \mathbb{k}e & \text{if } y = \chi(e)x, \\ 0 & \text{otherwise.} \end{cases}$$

The composition of morphisms is induced by the product of E . The monoidal product of $\mathbb{k}\mathcal{G}_\chi$ is the linear extension of the product of the groupoid \mathcal{G}_χ :

$$x \otimes y = xy \quad \text{and} \quad \left(x \xrightarrow{e} y\right) \otimes \left(z \xrightarrow{f} t\right) = \left(xz \xrightarrow{e^x f} yt\right).$$

The χ -category \mathcal{G}_χ is pivotal. The pivotal structures of \mathcal{G}_χ are parameterized by the group homomorphisms $d: H \rightarrow \mathbb{k}^*$. Given such a d , the dual of an object $x \in H$ is $x^* = x^{-1}$ with left and right (co)evaluations given by

$$\text{ev}_x = 1, \quad \text{coev}_x = 1, \quad \tilde{\text{ev}}_x = d(x)1, \quad \widetilde{\text{coev}}_x = d(x)^{-1}1,$$

where 1 is the unit element of E .

5.6. Particular cases

1. Given a group H , the trivial map $1 \rightarrow H$ is a crossed module and the notion of a $(1 \rightarrow H)$ -category agrees with that of an H -graded monoidal category.

2. Given an abelian group E , the trivial map $E \rightarrow 1$ is a crossed module and the notion of an $(E \rightarrow 1)$ -category agrees with that of a \mathbb{k} -linear monoidal category such that the Hom-sets are E -graded \mathbb{k} -modules, the composition and monoidal product of morphisms are E -multiplicative (i.e., multiplicative with respect to the degree of morphisms), and the identities and monoidal constraints are of degree $1 \in E$. In other words, $(E \rightarrow 1)$ -categories are monoidal categories enriched over the symmetric category of E -graded \mathbb{k} -modules.

6. Hopf group-coalgebras and their modules

Throughout this section, H denotes a group with neutral element 1. We review the definitions and basic properties of Hopf H -coalgebras and their modules (referring to [19] for details).

6.1. Group-coalgebras

An H -coalgebra (over \mathbb{k}) is a family of \mathbb{k} -modules $A = \{A_x\}_{x \in H}$ endowed with a family of \mathbb{k} -linear homomorphisms $\Delta = \{\Delta_{x,y}: A_{xy} \rightarrow A_x \otimes A_y\}_{x,y \in H}$ (the *coproduct*) and with a \mathbb{k} -linear homomorphism $\varepsilon: A_1 \rightarrow \mathbb{k}$ (the *counit*) which are coassociative and counital in the sense that for all $x, y, z \in H$,

$$(\Delta_{x,y} \otimes \text{id}_{A_z})\Delta_{xy,z} = (\text{id}_{A_x} \otimes \Delta_{y,z})\Delta_{x,yz}$$

and

$$(\text{id}_{A_x} \otimes \varepsilon)\Delta_{x,1} = \text{id}_{A_x} = (\varepsilon \otimes \text{id}_{A_x})\Delta_{1,x}.$$

Note that $(A_1, \Delta_{1,1}, \varepsilon)$ is a coalgebra (over \mathbb{k}) in the usual sense.

To any H -coalgebra $A = (\{A_x\}_{x \in H}, \Delta, \varepsilon)$ and any \mathbb{k} -algebra B , one associates the H -graded \mathbb{k} -algebra

$$\text{Conv}(A, B) = \bigoplus_{x \in H} \text{Hom}_{\mathbb{k}}(A_x, B),$$

called the *convolution algebra*, whose unit is $\varepsilon 1_B$ and whose product is defined by

$$f * g = m(f \otimes g)\Delta_{x,y} \in \text{Hom}_{\mathbb{k}}(A_{xy}, B)$$

for all $f \in \text{Hom}_{\mathbb{k}}(A_x, B)$ and $g \in \text{Hom}_{\mathbb{k}}(A_y, B)$, where m and 1_B are the product and unit of B .

6.2. Group-bicoalgebras

An H -bicoalgebra (over \mathbb{k}) is an H -coalgebra (over \mathbb{k}) $A = (\{A_x\}_{x \in H}, \Delta, \varepsilon)$ such that each A_x is endowed with a structure of a \mathbb{k} -algebra so that ε and $\Delta_{x,y}$ are algebra homomorphisms for all $x, y \in H$, that is,

$$\begin{aligned} \Delta_{x,y} \mu_{xy} &= (\mu_x \otimes \mu_y)(\text{id}_{A_x} \otimes \sigma_{A_y, A_x} \otimes \text{id}_{A_y})(\Delta_{x,y} \otimes \Delta_{x,y}), & \varepsilon \mu_1 &= \varepsilon \otimes \varepsilon, \\ \Delta_{x,y}(1_{xy}) &= 1_x \otimes 1_y, & \varepsilon(1_1) &= 1_{\mathbb{k}}, \end{aligned}$$

where $\mu_x: A_x \otimes A_x \rightarrow A_x$ and $1_x \in A_x$ denote the product and the unit element of A_x . Here and below, for \mathbb{k} -modules M and N , the flip $\sigma_{M,N}: M \otimes N \rightarrow N \otimes M$ is defined by $\sigma_{M,N}(u \otimes v) = v \otimes u$ for all $u \in M$ and $v \in N$.

6.3. Antipodes

An *antipode* for an H -bicoalgebra $A = (\{A_x\}_{x \in H}, \Delta, \varepsilon)$ is a family $S = \{S_x: A_{x^{-1}} \rightarrow A_x\}_{x \in H}$ of \mathbb{k} -linear homomorphisms such that for all $x \in H$,

$$\mu_x(S_x \otimes \text{id}_{A_x})\Delta_{x^{-1},x} = \eta_x \varepsilon = \mu_x(\text{id}_{A_x} \otimes S_x)\Delta_{x,x^{-1}}$$

where μ_x and $\eta_x = (1_{\mathbb{k}} \in \mathbb{k} \mapsto 1_x \in A_x)$ are the product and the unit map of A_x . This axiom means that for any $x \in H$, the homomorphism S_x is the inverse of id_{A_x} in the convolution algebra $\text{Conv}(A, A_x)$. An antipode $S = \{S_x\}_{x \in H}$ is *bijective* if S_x is bijective for all $x \in H$.

If it exists, an antipode is unique. Also, it is anti-multiplicative: for all $x \in H$,

$$S_x \mu_{x^{-1}} = \mu_x \sigma_{A_x, A_x}(S_x \otimes S_x) \quad \text{and} \quad S_x(1_{x^{-1}}) = 1_x,$$

and anti-comultiplicative: for all $x, y \in H$,

$$\Delta_{x,y}S_{xy} = (S_x \otimes S_y)\sigma_{A_{y^{-1}}, A_{x^{-1}}}\Delta_{y^{-1}, x^{-1}} \quad \text{and} \quad \varepsilon S_1 = \varepsilon.$$

6.4. Hopf group-coalgebras

A Hopf H -coalgebra is an H -bicoalgebra endowed with a bijective antipode. When $H = 1$, one recovers the usual notion of a Hopf algebra. In particular, if A is a Hopf H -coalgebra, then A_1 is a Hopf algebra.

The product μ_x , unit map η_x , coproduct $\Delta_{x,y}$, counit ε , antipode S_x and its inverse S_x^{-1} of a Hopf H -coalgebra $A = \{A_x\}_{x \in H}$ are depicted as follows:

$$\mu_x = \begin{array}{c} | \\ \text{---} \\ \bigcap \\ | \\ x \end{array} \quad \eta_x = \begin{array}{c} | \\ \bigcirc \\ | \\ x \end{array} \quad \Delta_{x,y} = \begin{array}{c} \bigcup \\ | \quad | \\ x \quad y \end{array} \quad \varepsilon = \begin{array}{c} \bigcirc \\ | \\ 1 \end{array} \quad S_x = \begin{array}{c} | \\ \oplus \\ | \\ x^{-1} \end{array} \quad S_x^{-1} = \begin{array}{c} | \\ \ominus \\ | \\ x \end{array}.$$

Here the colors $x, y \in H$ are abbreviations for A_x and A_y . The axioms of a Hopf H -coalgebra are then easily depicted in a manner similar to that of Sections 3.1-3.4.

An H -grouplike element of a Hopf H -coalgebra A is a family $G = (G_x)_{x \in H}$ with $G_x \in A_x$ such that for all $x, y \in H$,

$$\Delta_{x,y}(G_{xy}) = G_x \otimes G_y \quad \text{and} \quad \varepsilon(G_1) = 1_{\mathbb{k}}.$$

Note that each G_x is then invertible in A_x with inverse

$$G_x^{-1} = S_x(G_{x^{-1}}) = S_x^{-1}(G_{x^{-1}}),$$

where $S = \{S_x\}_{x \in H}$ is the antipode of A . The set $G_H(A)$ of H -grouplike elements of A is thus a group for the pointwise product.

6.5. Modules over Hopf group-coalgebras

Any H -bicoalgebra $A = \{A_x\}_{x \in H}$ (over \mathbb{k}) yields the H -graded \mathbb{k} -linear monoidal category $\text{Mod}_H(A)$ of A -modules and A -linear morphisms defined as follows.

A (left) A -module is an H -graded \mathbb{k} -module

$$M = \bigoplus_{x \in H} M_x$$

such that each M_x is endowed with a structure of a (left) module over the \mathbb{k} -algebra A_x .

An (H, A) -linear morphism between two A -modules M, N is a \mathbb{k} -linear homomorphism $\alpha: M \rightarrow N$ such that:

- (a) The map α preserves the H -grading: $\alpha(M_x) \subset N_x$ for all $x \in H$.
- (b) For all $x \in H$, the map $m \in M_x \mapsto \alpha(m) \in N_x$ is A_x -linear.


We let $\text{Mod}_H(A)$ be the category of A -modules and (H, A) -linear morphisms, with composition induced in the obvious way from the set-theoretical composition. The monoidal product of two A -modules M and N is the A -module

$$M \otimes N = \bigoplus_{x \in H} (M \otimes N)_x,$$

where the degree x summand

$$(M \otimes N)_x = \bigoplus_{\substack{y, z \in H \\ yz=x}} M_y \otimes N_z$$

is endowed with the A_x -action defined on $M_y \otimes N_z$ by



Here, the crossing denotes the symmetry (given by the usual flip maps) of the category $\text{Mod}_{\mathbb{k}}$ of \mathbb{k} -modules. The monoidal product of (H, A) -linear morphisms is their tensor product over \mathbb{k} . The unit object $\mathbb{1}$ is \mathbb{k} concentrated in degree $1 \in H$ with action given by the counit $\varepsilon: A_1 \rightarrow \mathbb{k}$. The monoidal constraints of $\text{Mod}_H(A)$ are inherited from the category $\text{Mod}_{\mathbb{k}}$. Then $\text{Mod}_H(A)$ is a \mathbb{k} -linear monoidal category. Note that the forgetful functor $\text{Mod}_H(A) \rightarrow \text{Mod}_{\mathbb{k}}$ is strict monoidal.

For any $x \in H$, by viewing any A_x -module as an A -module concentrated in degree x , the category $\text{Mod}_{\mathbb{k}}(A_x) = \text{Mod}_{\text{Mod}_{\mathbb{k}}}(A_x)$ of A_x -modules (see Section 3.6) is a full subcategory of $\text{Mod}_H(A)$. Then the category $\text{Mod}_H(A)$ is H -graded by the family $\{\text{Mod}_{\mathbb{k}}(A_x)\}_{x \in H}$:

$$\text{Mod}_H(A) = \bigoplus_{x \in H} \text{Mod}_{\mathbb{k}}(A_x)$$

In particular, the homogeneous objects of $\text{Mod}_H(A)$ are the nonzero A_x -modules where x runs over H . Note that $\text{Mod}_H(A)$ is additive: any finite direct sum of A -modules always exists (it is induced in the obvious way from the direct sums in $\text{Mod}_{\mathbb{k}}$). Furthermore $\text{Mod}_H(A)$ is abelian (with kernels and cokernels induced in the obvious way from those in $\text{Mod}_{\mathbb{k}}$).

Let $\text{mod}_H(A)$ be the full subcategory of $\text{Mod}_H(A)$ consisting of the A -modules whose underlying \mathbb{k} -module is projective of finite rank. Then $\text{mod}_H(A)$ is a \mathbb{k} -linear monoidal subcategory of $\text{Mod}_H(A)$. It is additive and H -graded:

$$\text{mod}_H(A) = \bigoplus_{x \in H} \text{mod}_{\mathbb{k}}(A_x).$$

where $\text{mod}_{\mathbb{k}}(A_x)$ is the full subcategory of $\text{Mod}_{\mathbb{k}}(A_x)$ consisting of the A_x -modules whose underlying \mathbb{k} -module is projective of finite rank.

If A is a Hopf H -coalgebra, then the monoidal category $\text{Mod}_H(A)$ is closed (so that the forgetful functor $\text{Mod}_H(A) \rightarrow \text{Mod}_{\mathbb{k}}$ preserves the internal Homs) and the monoidal category $\text{mod}_H(A)$ is rigid. The formulas for the internal Homs and the duals of objects are similar (with obvious changes) to those given in Section 3.6.

Assume that A is a Hopf H -coalgebra. A *pivotal element* for A is an H -grouplike element $G = (G_x)_{x \in H}$ of A (see Section 6.4) such that for all $x \in H$ and $a \in A_x$,

$$S_x S_{x^{-1}}(a) = G_x a G_x^{-1}.$$

Then pivotal structures on $\text{mod}_H(A)$ are in bijective correspondence with pivotal elements of A . The pivotal structure on $\text{mod}_H(A)$ associated with a pivotal element $G = (G_x)_{x \in H}$ is given as follows: the dual of an

object $(M, r) \in \text{mod}_{\mathbb{k}}(A_x)$ is the object $(M, r)^* = (M^*, r^\dagger) \in \text{mod}_{\mathbb{k}}(A_{x^{-1}})$, where $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$ and the action r^\dagger is given as in Section 3.6: for all $a \in A_{x^{-1}}$, $\varphi \in M^*$, and $m \in M$,

$$r^\dagger(a \otimes \varphi)(m) = \varphi(S_x(am)).$$

The associated (co)evaluations are given, for all $\varphi \in M^*$, $m \in M$, by

$$\begin{aligned} \text{ev}_{(M,r)}(\varphi \otimes m) &= \varphi(m), & \tilde{\text{ev}}_M(m \otimes \varphi) &= \varphi(G_x m), \\ \text{coev}_{(M,r)}(1_{\mathbb{k}}) &= \sum_i b_i \otimes b_i^*, & \widetilde{\text{coev}}_{(M,r)}(1_{\mathbb{k}}) &= \sum_i b_i^* \otimes G_x^{-1} b_i, \end{aligned}$$

where $(b_i)_i$ is any basis of M and $(b_i^*)_i$ is the dual basis of M^* .

6.6. Example

Consider the trivial Hopf H -coalgebra $\mathbb{k}_H = \{(\mathbb{k}_H)_x = \mathbb{k}\}_{x \in H}$ whose structural morphisms are given for all $x, y \in H$ by

$$\Delta_{x,y}(1_{\mathbb{k}}) = 1_{\mathbb{k}} \otimes 1_{\mathbb{k}}, \quad \varepsilon = \text{id}_{\mathbb{k}}, \quad S_x = \text{id}_{\mathbb{k}}.$$

Then the closed H -graded monoidal category $\text{Mod}_H(\mathbb{k}_H)$ is nothing but the category of H -graded \mathbb{k} -modules and \mathbb{k} -linear grading-preserving homomorphisms.

6.7. Remark

A homomorphism from an H -bicoalgebra $A = \{A_x\}_{x \in H}$ to an H -bicoalgebra $B = \{B_x\}_{x \in H}$ is a family $f = \{f_x: A_x \rightarrow B_x\}_{x \in H}$ of algebra homomorphisms compatible with the coproducts and counits of A and B in the following sense: for all $x, y \in H$,

$$\Delta_{x,y}^B f_{xy} = (f_x \otimes f_y) \Delta_{x,y}^A \quad \text{and} \quad \varepsilon^B f_1 = \varepsilon^A.$$

Note that if A and B are Hopf H -coalgebras, then any H -bicoalgebra homomorphism $f: A \rightarrow B$ preserves the antipodes of A and B , that is, $S_x^B f_{x^{-1}} = f_x S_x^A$ for all $x \in H$.

It is not difficult to check that any H -bicoalgebra homomorphism $f: A \rightarrow B$ induces an H -graded functor $f^*: \text{Mod}_H(B) \rightarrow \text{Mod}_H(A)$ that is, a strong monoidal \mathbb{k} -linear functor that preserves the H -grading of objects. Moreover, if A and B are Hopf H -coalgebras, then the functor f^* is closed, that is, it preserves the internal Homs. Also, if A and B are Hopf H -coalgebras endowed with pivotal elements, then any H -bicoalgebra homomorphism $f: A \rightarrow B$ preserving the pivotal elements of A and B (that is, $f_x(G_x^A) = G_x^B$ for all $x \in H$) induces a pivotal H -graded functor $f^*: \text{mod}_H(B) \rightarrow \text{mod}_H(A)$.

7. Hopf crossed module-coalgebras

Throughout this section, $\chi: E \rightarrow H$ is a crossed module. We introduce the notions of χ -bicoalgebras and Hopf χ -coalgebras. The main motivation is that their categories of representations (introduced in Section 8) are χ -graded monoidal categories.

7.1. Crossed module-actions

A χ -action on an H -bicoalgebra $A = \{A_x\}_{x \in H}$ is a family of \mathbb{k} -algebra homomorphisms

$$\phi = \{\phi_{x,e} : A_x \rightarrow A_{\chi(e)x}\}_{(x,e) \in H \times E}$$

such that for all $x, y \in H$ and $e, f \in E$,

$$\phi_{x,1} = \text{id}_{A_x}, \tag{2}$$

$$\phi_{\chi(e)x,f} \phi_{x,e} = \phi_{x,fe}, \tag{3}$$


$$(\phi_{x,e} \otimes \phi_{y,f}) \Delta_{x,y} = \Delta_{\chi(e)x, \chi(f)y} \phi_{xy, e^x f}. \tag{4}$$

Note the indices in Axiom (4) are coherent since the equivariance of χ (see Section 4.1) implies that $\chi(e^x f)xy = \chi(e)x\chi(f)y$. Also, it follows directly from (2) and (3) that each $\phi_{x,e}$ is an isomorphism and


$$\phi_{x,e}^{-1} = \phi_{\chi(e)x, e^{-1}}.$$

A χ -action is *trivial* if $\phi_{x,e} = \text{id}_{A_x}$ (and so $A_{\chi(e)x} = A_x$) for all $x \in H$ and $e \in E$.


We depict the χ -action $\phi_{x,e} : A_x \rightarrow A_{\chi(e)x}$ by a strand with a dot labeled with e (on the left or on the right) as follows:



Axioms (2)-(4) are depicted as



The fact that $\phi_{x,e}$ is an algebra homomorphism is depicted as



7.2. Hopf crossed module-coalgebras

A χ -bicoalgebra (over \mathbb{k}) is an H -bicoalgebra (over \mathbb{k}) endowed with a χ -action.

A Hopf χ -coalgebra (over \mathbb{k}) is a Hopf H -coalgebra (over \mathbb{k}) endowed with a χ -action. Equivalently, a Hopf χ -coalgebra is a χ -bicoalgebra whose underlying H -bicoalgebra has a bijective antipode.

In Section 10, we prove that Hopf crossed module-coalgebras may be seen as Hopf algebras in some symmetric monoidal category (see Corollary 10.3). In the next lemma, we show that the antipode of a Hopf χ -coalgebra is compatible with the χ -action.

Lemma 7.1. *Let $A = \{A_x\}_{x \in H}$ be a Hopf χ -coalgebra, with antipode $S = \{S_x\}_{x \in H}$ and χ -action $\phi = \{\phi_{x,e}\}_{(x,e) \in H \times E}$. Then*

$$\phi_{x,e} S_x = S_{\chi(e)x} \phi_{x^{-1}, x^{-1}(e^{-1})}$$


for all $x \in H$ and $e \in E$.

Proof. Axiom (4) and the multiplicativity of $\phi_{x,e}^{-1} = \phi_{\chi(e)x, e^{-1}}$ imply that

$$\phi_{x,e}^{-1} S_{\chi(e)x} \phi_{x^{-1}, x^{-1}(e^{-1})} : A_{x^{-1}} \rightarrow A_x$$

is inverse to id_{A_x} in the convolution algebra $\text{Conv}(A, A_x)$, and so is equal to S_x . \square

Graphically, the compatibility of the antipode with the χ -action (see Lemma 7.1) is depicted as



7.3. Particular cases

1. Given a group H , the trivial map $1 \rightarrow H$ is a crossed module and it follows from (2) that any $(1 \rightarrow H)$ -action for an H -bicoalgebra is trivial. Consequently, the notion of a Hopf $(1 \rightarrow H)$ -coalgebra agrees with that of a Hopf H -coalgebra.

2. Given an abelian group E , the trivial map $E \rightarrow 1$ is a crossed module and the notion of a Hopf $(E \rightarrow 1)$ -coalgebra agrees with that of a Hopf algebra A endowed with an action of E by algebra and bicomodule automorphisms, that is, with a group homomorphism $\phi: E \rightarrow \text{Aut}_{\mathbb{k}}(A)$ such that for all $e \in E$,

- ϕ_e is an algebra automorphism of A ,
- ϕ_e is a bicomodule automorphism of A , that is, $\Delta \phi_e = (\phi_e \otimes \text{id}_A)\Delta = (\text{id}_A \otimes \phi_e)\Delta$.

Here $\text{Aut}_{\mathbb{k}}(A)$ denotes the group of \mathbb{k} -linear automorphisms of A and Δ is the coproduct of A .

7.4. Example

Let $\omega: E \times G \rightarrow \mathbb{k}^*$ be a bicharacter, where E is an abelian group and G is a group. Recall that the group algebra $\mathbb{k}[G]$ is a Hopf algebra with coproduct Δ , counit ε , and antipode S defined by $\Delta(g) = g \otimes g$, $\varepsilon(g) = 1_{\mathbb{k}}$, and $S(g) = g^{-1}$ for all $g \in G$. Consider the group homomorphism $\phi: E \rightarrow \text{Aut}_{\mathbb{k}}(\mathbb{k}[G])$ defined by $\phi_e(g) = \omega(e, g)g$ for all $e \in E$ and $g \in G$. Then $\mathbb{k}[G]$ is a Hopf $(E \rightarrow 1)$ -coalgebra with $(E \rightarrow 1)$ -action ϕ , which we denote by $\mathbb{k}^{\omega}[G]$.

7.5. Example

Let $\chi: E \rightarrow H$ be a crossed module, A be a Hopf \mathbb{k} -algebra, and $\rho: H \rightarrow \text{Aut}_{\text{Alg}}(A)$ be a group homomorphism, where $\text{Aut}_{\text{Alg}}(A)$ is the group of algebra automorphisms of A . Let δ , ε , and s be the coproduct, counit, and antipode of A , respectively. For any $x \in H$, set $A_x = A$ as a \mathbb{k} -algebra. Then the

family $A_\chi^c = \{A_x\}_{x \in H}$ is a Hopf χ -coalgebra with counit $\varepsilon: A_1 \rightarrow \mathbb{k}$ and with coproduct, antipode, and χ -action respectively defined by:

$$\begin{aligned} \Delta &= \{\Delta_{x,y} = (\rho_x \otimes \rho_y)\delta\rho_{(xy)^{-1}}: A_{xy} \rightarrow A_x \otimes A_y\}_{x,y \in H}, \\ S &= \{S_x = \rho_x s \rho_x: A_{x^{-1}} \rightarrow A_x\}_{x \in H}, \\ \phi &= \{\phi_{x,e} = \rho_{\chi(e)}: A_x \rightarrow A_{\chi(e)x}\}_{(x,e) \in H \times E}, \end{aligned}$$

where ρ_z denotes the image of $z \in H$ under ρ .

7.6. Example

Hopf χ -coalgebras with trivial χ -action are in bijective correspondence with Hopf π -coalgebras, where $\pi = \text{Coker}(\chi) = H/\text{Im}(\chi)$ is the so-called fundamental group of χ . Indeed, let $B = \{B_g\}_{g \in \pi}$ be a Hopf π -coalgebra. Denote by $p: H \rightarrow \pi$ the canonical projection. For any $x \in H$, set $A_x = B_{p(x)}$ as a \mathbb{k} -algebra. For any $x, y \in H$, set $\Delta_{x,y} = \delta_{p(x),p(y)}$ and $S_x = s_{p(x)}$, where δ and s are the coproduct and antipode of B . Then the family $A = \{A_x\}_{x \in H}$, endowed with the coproduct Δ , the same counit of B , and the antipode S , is a Hopf χ -coalgebra with trivial χ -action.

Conversely, let $A = \{A_x\}_{x \in H}$ be a Hopf χ -coalgebra with trivial χ -action. Pick a set-theoretical section $q: \pi \rightarrow H$ of the canonical projection $p: H \rightarrow \pi$, meaning that $pq = \text{id}_\pi$. Note that the triviality of the χ -action implies that $A_x = A_y$ for all $x, y \in H$ such that $p(x) = p(y)$. For any $\alpha, \beta \in \pi$, set $B_\alpha = A_{q(\alpha)}$ as a \mathbb{k} -algebra, $\delta_{\alpha,\beta} = \Delta_{q(\alpha),q(\beta)}$, and $s_\alpha = S_{q(\alpha)}$, where Δ and S are the coproduct and antipode of A . Then the family $B = \{B_\alpha\}_{\alpha \in \pi}$, endowed with the coproduct δ , the same counit of A , and the antipode s , is a Hopf π -coalgebra. Note that B does not depend on the choice of the section q because the χ -action of A is trivial.

7.7. Grouplike elements

An H -grouplike element of a Hopf χ -coalgebra A is an H -grouplike element the Hopf H -coalgebra underlying A . As in Section 6.4, we denote by $G_H(A)$ the group of H -grouplike elements of A .

A χ -grouplike element of a Hopf χ -coalgebra A is an H -grouplike element $G = (G_x)_{x \in H}$ of A which is χ -equivariant in the sense that for all $x \in H$ and $e \in E$,

$$\phi_{x,e}(G_x) = G_{\chi(e)x}.$$

The set $G_\chi(A)$ of χ -grouplike elements of A is a subgroup of $G_H(A)$. Note that it may be a strict subgroup. For example, for the Hopf $(E \rightarrow 1)$ -coalgebra $\mathbb{k}^\omega[G]$ of Example 7.4, we have:

$$G_1(\mathbb{k}^\omega[G]) = G \quad \text{and} \quad G_{E \rightarrow 1}(\mathbb{k}^\omega[G]) = \{g \in G \mid \omega(e, g) = 1_{\mathbb{k}} \text{ for all } e \in E\}.$$

The next lemma describes the behavior of the χ -action on H -grouplike elements.

Lemma 7.2. *Let A be a Hopf χ -coalgebra. Then the map*

$$G_H(A) \times E \rightarrow \mathbb{k}^*, \quad (G, e) \mapsto \langle G, e \rangle = \varepsilon(\phi_{\chi(e^{-1}),e}(G_{\chi(e^{-1})}))$$

is a bicharacter such that for all $G \in G_H(A)$ and $(x, e) \in H \times E$,

$$\phi_{x,e}(G_x) = \langle G, e \rangle G_{\chi(e)x}.$$

Here, ε is the counit and ϕ is the χ -action of A .

It follows directly from Lemma 7.2 that an H -grouplike element G of A is a χ -grouplike element if and only if $\langle G, e \rangle = 1$ for all $e \in E$.

Proof. For all $G \in G_H(A)$ and $(x, e) \in H \times E$, we have:

$$\begin{aligned} \phi_{x,e}(G_x) &\stackrel{(i)}{=} (\varepsilon \otimes \text{id}_{A_{\chi(e)x}})\Delta_{1,\chi(e)x} \phi_{x,e}(G_x) \\ &\stackrel{(ii)}{=} (\varepsilon \phi_{\chi(e^{-1}),e} \otimes \phi_{\chi(e)x,1})\Delta_{\chi(e^{-1}),\chi(e)x}(G_x) \\ &\stackrel{(iii)}{=} \varepsilon(\phi_{\chi(e^{-1}),e}(G_{\chi(e^{-1})})) \phi_{\chi(e)x,1}(G_{\chi(e)x}) \\ &\stackrel{(iv)}{=} \langle G, e \rangle G_{\chi(e)x}. \end{aligned}$$

Here (i) follows the counitality of the coproduct, (ii) from (4), (iii) from the fact that G is H -grouplike, and (iv) from (2) and the definition of $\langle G, e \rangle$. The multiplicativity of the map $(G, e) \mapsto \langle G, e \rangle$ in the variable G follows from the multiplicativity of the counit and of the maps $\phi_{\chi(e^{-1}),e}$. Its multiplicativity in the variable e follows from (4). Hence this map is indeed a bicharacter $G_H(A) \times E \rightarrow \mathbb{k}^*$. \square

7.8. The dual notion: Hopf crossed module-algebras

The notion of a Hopf χ -coalgebra is not self dual. Its dual notion is that of a Hopf χ -algebra defined as follows. Recall that a \mathbb{k} -algebra A is H -graded if it is endowed with a direct sum decomposition

$$A = \bigoplus_{x \in H} A_x$$

such that $1_A \in A_1$ and $A_x A_y \subset A_{xy}$ for all $x, y \in H$.

An H -bialgebra (over \mathbb{k}) is an H -graded \mathbb{k} -algebra $A = \bigoplus_{x \in H} A_x$ such that each A_x is endowed with a structure of a \mathbb{k} -coalgebra so that the unit map $\eta: \mathbb{k} \rightarrow A$ (defined by $\eta(1_{\mathbb{k}}) = 1_A$) and the restricted products $\mu_{x,y}: A_x \otimes A_y \rightarrow A_{xy}$ are coalgebra homomorphisms for all $x, y \in H$, that is,

$$\begin{aligned} \Delta_{xy} \mu_{x,y} &= (\mu_{x,y} \otimes \mu_{x,y})(\text{id}_{A_x} \otimes \sigma_{A_x, A_y} \otimes \text{id}_{A_y})(\Delta_x \otimes \Delta_y), & \varepsilon_{xy} \mu_{x,y} &= \varepsilon_x \otimes \varepsilon_y, \\ \Delta_1(1_A) &= 1_A \otimes 1_A, & \varepsilon_1(1_A) &= 1_{\mathbb{k}}, \end{aligned}$$

where $\Delta_x: A_x \rightarrow A_x \otimes A_x$ and $\varepsilon_x: A_x \rightarrow \mathbb{k}$ denote the coproduct and counit of A_x , and σ_{A_x, A_y} is the usual flip.

A Hopf H -algebra is an H -bialgebra $A = \bigoplus_{x \in H} A_x$ endowed with a bijective antipode, that is, a family $S = \{S_x: A_x \rightarrow A_{x^{-1}}\}_{x \in H}$ of \mathbb{k} -linear isomorphisms such that for all $x \in H$,

$$\mu_{x^{-1},x}(S_x \otimes \text{id}_{A_x})\Delta_x = \varepsilon_x 1_A = \mu_{x,x^{-1}}(\text{id}_{A_x} \otimes S_x)\Delta_x.$$

A χ -action on a H -bialgebra $A = \bigoplus_{x \in H} A_x$ is a family

$$\phi = \{\phi_{x,e}: A_x \rightarrow A_{\chi(e)x}\}_{x \in H, e \in E}$$

of \mathbb{k} -coalgebra homomorphisms such that for all $x, y \in H$ and $e, f \in E$,

- $\phi_{x,1} = \text{id}_{A_x}$,
- $\phi_{\chi(e)x,f} \phi_{x,e} = \phi_{x,fe}$,
- $\mu_{\chi(e)x,\chi(f)y}(\phi_{x,e} \otimes \phi_{y,f}) = \phi_{xy,e^x f} \mu_{x,y}$.

Note that each $\phi_{x,e}$ is then an isomorphism and $\phi_{x,e}^{-1} = \phi_{\chi(e)x,e^{-1}}$.

A χ -bialgebra is an H -bialgebra endowed with a χ -action. A Hopf χ -algebra is a Hopf H -algebra endowed with a χ -action.

A Hopf χ -(co)algebra A is of *finite type* if for all $x \in H$, the \mathbb{k} -module A_x is projective of finite rank.

The dual of a Hopf χ -algebra of finite type is a Hopf χ -coalgebra (of finite type). More explicitly, let $A = \bigoplus_{x \in H} A_x$ be a Hopf χ -algebra of finite type. For any \mathbb{k} -module M , set $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$. Then the dual $A^* = \{A_x^*\}_{x \in H}$ of A is a Hopf χ -coalgebra. Its coproduct $\Delta_{x,y}: A_{xy}^* \rightarrow A_x^* \otimes A_y^*$ is the composition of the transpose of the restricted product $A_x \otimes A_y \rightarrow A_{xy}$ with the canonical isomorphism $(A_x \otimes A_y)^* \cong A_x^* \otimes A_y^*$. The algebra structure of A^* is induced in the standard way by the coalgebra structure of A_x . The antipode and χ -action of A^* are the transpose of those of A .

Conversely, the dual $A^* = \bigoplus_{x \in H} A_x^*$ of a Hopf χ -coalgebra $A = \{A_x\}_{x \in H}$ of finite type is a Hopf χ -algebra (with transposed structural morphisms).

8. Categories of representations of Hopf crossed module-coalgebras

Throughout this section, $\chi: E \rightarrow H$ is a crossed module and $A = \{A_x\}_{x \in H}$ is a χ -bicoalgebra (over \mathbb{k}). We associate to A two categories of representations $\text{Mod}_{\chi}(A)$ and $\text{mod}_{\chi}(A)$ which are χ -graded categories (in the sense of Section 5.4).

8.1. Modules over χ -bicoalgebras

A (left) A -module is a module over the H -bicoalgebra underlying A , that is, an H -graded \mathbb{k} -module

$$M = \bigoplus_{x \in H} M_x$$

such that each M_x is endowed with a structure of an A_x -module (see Section 6.5).

A (χ, A) -linear morphism between two A -modules M and N is a \mathbb{k} -linear homomorphism $\alpha: M \rightarrow N$ such that:


- (a) For all $x \in H$,

$$\alpha(M_x) \subset \bigoplus_{e \in E} N_{\chi(e)x}.$$

- (b) For all $x \in H$ and $e \in E$, the \mathbb{k} -linear homomorphism $\alpha_{x,e}: M_x \rightarrow N_{\chi(e)x}$ induced by α (restriction to M_x followed by projection to $N_{\chi(e)x}$) is an A_x -linear morphism

$$\alpha_{x,e}: M_x \rightarrow \phi_{x,e}^*(N_{\chi(e)x}),$$

where the A_x -module $\phi_{x,e}^*(N_{\chi(e)x})$ is the pullback of the $A_{\chi(e)x}$ -module $N_{\chi(e)x}$ along the algebra isomorphism $\phi_{x,e}: A_x \rightarrow A_{\chi(e)x}$ given by the χ -action of A . Graphically, the A_x -linearity of $\alpha_{x,e}$ depicts as:



8.2. The categories of representations

We let $\text{Mod}_\chi(A)$ be the category of A -modules and (χ, A) -linear morphisms, with composition induced in the obvious way from the set-theoretical composition. The category $\text{Mod}_\chi(A)$ is \mathbb{k} -linear (as a subcategory of $\text{Mod}_{\mathbb{k}}$). For any two A -modules M and N , we endow the \mathbb{k} -module $\text{Hom}_{\text{Mod}_\chi(A)}(M, N)$ with a structure of an E -graded \mathbb{k} -module by setting:

$$\text{Hom}_{\text{Mod}_\chi(A)}(M, N) = \bigoplus_{e \in E} \text{Hom}_{\text{Mod}_\chi(A)}^e(M, N)$$

where $\text{Hom}_{\text{Mod}_\chi(A)}^e(M, N)$ is the set of (χ, A) -linear morphisms $\alpha: M \rightarrow N$ such that for all $x \in E$,

$$\alpha(M_x) \subset N_{\chi(e)x}.$$

The axioms (2) and (3) of a χ -action imply that $\text{Mod}_\chi(A)$ is E -Hom-graded. Also, since the (χ, A) -linear morphisms of degree 1 are nothing but the (H, A) -linear morphisms, the 1-subcategory of $\text{Mod}_\chi(A)$ is

$$\text{Mod}_\chi(A)^1 = \text{Mod}_H(A),$$

where $\text{Mod}_H(A)$ is the category of modules over the H -bicoalgebra underlying A (see Section 6.5). Note that for all A -modules M, N and $e \in E$,

$$\text{Hom}_{\text{Mod}_\chi(A)}^e(M, N) = \text{Hom}_{\text{Mod}_H(A)}(M, \phi_e^*(N)),$$

where $\phi_e^*(N)$ is the A -module defined by

$$\phi_e^*(N) = \bigoplus_{x \in H} \phi_e^*(N)_x \quad \text{with} \quad \phi_e^*(N)_x = \phi_{x,e}^*(N_{\chi(e)x}) \in \text{Mod}_{\mathbb{k}}(A_x).$$

The category $\text{Mod}_\chi(A)$ is monoidal: the monoidal product of two A -modules, the unit object, and the monoidal constraints of $\text{Mod}_\chi(A)$ are those of $\text{Mod}_H(A)$, and the monoidal product of two (χ, A) -linear morphisms is their tensor product over \mathbb{k} .

Theorem 8.1. *The category $\text{Mod}_\chi(A)$ is a χ -graded monoidal category which is E -additive (see Section 5.3) and abelian. Moreover, if A is a Hopf χ -coalgebra, then $\text{Mod}_\chi(A)$ is closed.*

Proof. By the above, the category $\text{Mod}_\chi(A)$ is E -Hom-graded and its 1-subcategory is $\text{Mod}_\chi(A)^1 = \text{Mod}_H(A)$. In particular, $\text{Mod}_\chi(A)^1$ is H -graded (see Section 6.5). The fact that the tensor product over \mathbb{k} of two (χ, A) -linear morphisms is a (χ, A) -linear morphism follows from (4). This implies that the above definitions do define a monoidal structure on $\text{Mod}_\chi(A)$.

To prove that $\text{Mod}_\chi(A)$ is E -additive, first remark that finite 1-direct sums of A -modules exist in $\text{Mod}_\chi(A)$ (since $\text{Mod}_H(A)$ is additive by Section 6.5). Then, for any $e \in E$, the e -direct sum of a finite family $(M_\lambda)_{\lambda \in \Lambda}$ of A -modules exists in $\text{Mod}_\chi(A)$ and is computed by

$$\bigoplus_{\lambda \in \Lambda}^e M_\lambda = \bigoplus_{\lambda \in \Lambda}^1 \phi_{e^{-1}}^*(M_\lambda).$$

The fact that $\text{Mod}_\chi(A)$ is abelian is proved as in the classical case by noticing that the kernels and cokernels in $\text{Mod}_\chi(A)$ are induced in the obvious way from those in $\text{Mod}_{\mathbb{k}}$.

Finally, if A is a Hopf χ -coalgebra, then $\text{Mod}_\chi(A)$ is closed since its 1-subcategory $\text{Mod}_\chi(A)^1 = \text{Mod}_H(A)$ is closed (see Section 6.5). \square

Let $\text{mod}_\chi(A)$ be the full subcategory of $\text{Mod}_\chi(A)$ consisting of the A -modules whose underlying \mathbb{k} -module is projective of finite rank. By a *pivotal element* of a Hopf χ -coalgebra, we mean a pivotal element of its underlying Hopf H -coalgebra (see Section 6.5).

Corollary 8.2. *The category $\text{mod}_\chi(A)$ is an E -additive χ -graded monoidal category. Moreover, if A is a Hopf χ -coalgebra, then $\text{mod}_\chi(A)$ is rigid and the pivotal structures on $\text{mod}_\chi(A)$ are in bijective correspondence with the pivotal elements of A .*

Note as above that the 1-subcategory of $\text{mod}_\chi(A)$ is

$$\text{mod}_\chi(A)^1 = \text{mod}_H(A),$$

where $\text{mod}_H(A)$ is the H -graded monoidal category associated with the Hopf H -coalgebra underlying A (see Section 6.5).

Proof. The first assertion follows from the fact that the monoidal structure of $\text{Mod}_\chi(A)$ restricts to $\text{mod}_\chi(A)$. Assume A is a Hopf χ -coalgebra. Then $\text{mod}_\chi(A)$ is rigid since its 1-subcategory $\text{mod}_\chi(A)^1 = \text{mod}_H(A)$ is rigid (see Section 6.5). Also, the pivotal structures on $\text{mod}_\chi(A)$ are in bijective correspondence with the pivotal structures on $\text{mod}_\chi(A)^1$, and so in bijective correspondence with the pivotal elements of A (see Section 6.5). \square

8.3. Remark

A homomorphism from a χ -bicoalgebra A to a χ -bicoalgebra B is an H -bicoalgebra homomorphism $f: A \rightarrow B$ (see Remark 6.7) which is compatible with the χ -actions of A and B in the following sense: for all $(x, e) \in H \times E$,

$$\phi_{x,e}^B f_x = f_{\chi(e)x} \phi_{x,e}^A.$$

It is not difficult to check that such a homomorphism $f: A \rightarrow B$ induces χ -graded functors

$$f^*: \text{Mod}_\chi(B) \rightarrow \text{Mod}_\chi(A) \quad \text{and} \quad f_*: \text{mod}_\chi(B) \rightarrow \text{mod}_\chi(A).$$

Here, by a χ -graded functor between χ -graded monoidal categories, we mean a strong monoidal \mathbb{k} -linear functor that preserves the H -grading of objects and the E -grading of morphisms.

8.4. Example

Consider the trivial Hopf χ -coalgebra $\mathbb{k}_\chi = \{(\mathbb{k}_\chi)_x = \mathbb{k}\}_{x \in H}$ whose structural morphisms are given for all $x, y \in H$ and $e \in E$ by

$$\Delta_{x,y}(\mathbb{1}_\mathbb{k}) = \mathbb{1}_\mathbb{k} \otimes \mathbb{1}_\mathbb{k}, \quad \varepsilon = \text{id}_\mathbb{k}, \quad S_x = \text{id}_\mathbb{k}, \quad \phi_{x,e} = \text{id}_\mathbb{k}.$$

By Theorem 8.1, the category $\text{Mod}_\chi(\mathbb{k}_\chi)$ is an E -additive closed χ -graded monoidal category. Its objects are the H -graded \mathbb{k} -modules $M = \bigoplus_{x \in H} M_x$ and its Hom-sets are computed by

$$\text{Hom}_{\text{Mod}_\chi(\mathbb{k}_\chi)}^e(M, N) = \bigoplus_{x \in H} \text{Hom}_\mathbb{k}(M_x, N_{\chi(e)x}).$$

The composition is induced from the set-theoretical composition. The monoidal product of objects is that of H -graded modules, the monoidal product of morphisms is their tensor product over \mathbb{k} , and the monoidal constraints are inherited from the category of \mathbb{k} -modules. Note that the 1-subcategory of $\text{Mod}_\chi(\mathbb{k}_\chi)$ is the category of H -graded \mathbb{k} -modules and \mathbb{k} -linear grading-preserving homomorphisms (see Example 6.6). Also, the χ -category $\mathbb{k}\mathcal{G}_\chi$ from Example 5.5 is isomorphic to the full subcategory of $\text{Mod}_\chi(\mathbb{k}_\chi)$ whose set of objects is $\{\mathbb{k}_x\}_{x \in H}$, where \mathbb{k}_x is \mathbb{k} concentrated in degree x .

8.5. Example

Consider the Hopf $(E \rightarrow 1)$ -coalgebra $\mathbb{k}^\omega[G]$ from Example 7.4, where E is an abelian group, G is a group, and $\omega: E \times G \rightarrow \mathbb{k}^*$ is a bicharacter. By Theorem 8.1, the category $\mathcal{C}_\omega = \text{Mod}_{(E \rightarrow 1)}(\mathbb{k}^\omega[G])$ is an E -additive closed $(E \rightarrow 1)$ -graded monoidal category. Its objects are the \mathbb{k} -linear representations of G . For any representations M, N of G and any $e \in E$, the \mathbb{k} -module $\text{Hom}_{\mathcal{C}_\omega}^e(M, N)$ is the set of \mathbb{k} -linear maps $\alpha: M \rightarrow N$ such that for all $g \in G$ and $m \in M$,

$$\alpha(g \cdot m) = \omega(e, g) (g \cdot \alpha(m)).$$

The composition in \mathcal{C}_ω is induced from the set-theoretical composition and the monoidal product in \mathcal{C}_ω is the usual tensor product of representations of G . Note that these are E -multiplicative as expected (see Section 5.6).

8.6. Example

Consider the Hopf χ -coalgebra A_ρ^χ from Example 7.5, where A is a Hopf algebra and $\rho: H \rightarrow \text{Aut}_{\text{Alg}}(A)$ is a group homomorphism. By Theorem 8.1, the category $\text{Mod}_\chi(A_\rho^\chi)$ is an E -additive closed χ -graded monoidal category. Its objects are the H -graded A -modules, that is, the A -modules M endowed with an H -grading $M = \bigoplus_{x \in H} M_x$ so that $a \cdot M_x \subset M_x$ for all $a \in A$ and $x \in H$. The Hom-sets are given by

$$\text{Hom}_{\text{Mod}_\chi(\mathbb{k}_\chi)}^e(M, N) = \bigoplus_{x \in H} \text{Hom}_{\text{Mod}_k(A)}(M_x, \rho_{\chi(e)}^*(N_{\chi(e)x})).$$

The composition is induced from the set-theoretical composition. The monoidal product is induced from the usual monoidal product of A -modules (using the coproduct of A) and of H -graded \mathbb{k} -modules. Note that when $A = \mathbb{k}$ we recover Example 8.4 (since ρ becomes trivial and $\mathbb{k}_\rho^\chi = \mathbb{k}_\chi$ as Hopf χ -coalgebras).

8.7. χ -Fusion categories from Hopf χ -coalgebras

We first recall the definition of a χ -fusion category from [14]. By a *simple* object of a \mathbb{k} -linear category, we mean an object whose set of endomorphisms is a free \mathbb{k} -module of rank 1. Note that if \mathcal{C} is a χ -category over \mathbb{k} , then a simple object of the 1-subcategory \mathcal{C}^1 of \mathcal{C} is nothing but an object i of \mathcal{C} such that $\text{End}_{\mathcal{C}}^1(i) = \mathbb{k} \text{id}_i$.

A χ -fusion category (over \mathbb{k}) is a rigid χ -category \mathcal{C} (over \mathbb{k}) such that:

- (a) The 1-subcategory \mathcal{C}^1 of \mathcal{C} is H -fusion, that is:
 - $\text{Hom}_{\mathcal{C}^1}(i, j) = 0$ whenever i, j are non-isomorphic simple objects of \mathcal{C}^1 ,
 - each object of \mathcal{C}^1 is a (finite) direct sum of simple objects of \mathcal{C}^1 ,
 - the unit object $\mathbb{1}$ is a simple object of \mathcal{C}^1 ,
 - for any $h \in H$, there are at least one and only finitely many (up to isomorphism in \mathcal{C}^1) homogeneous simple objects of \mathcal{C}^1 with degree h .

(b) For any $e \in E$, each object of \mathcal{C} is an e -direct sum of a finite family of simple objects of \mathcal{C}^1 .

For example, the χ -category $\mathbb{k}\mathcal{G}_\chi$ from Example 5.5 is χ -fusion.

Clearly, the Hom-sets in a χ -fusion category are free \mathbb{k} -modules of finite rank. Note that a χ -fusion category \mathcal{C} may not be semisimple (as a \mathbb{k} -linear category) while its 1-subcategory \mathcal{C}^1 always is. Indeed, a simple object i of \mathcal{C}^1 is not necessarily simple in \mathcal{C} : it may happen that $\text{End}_{\mathcal{C}}^e(i) \neq 0$ for some $e \in E \setminus \{1\}$ (as for example in $\mathbb{k}\mathcal{G}_\chi$ when $\text{Ker}(\chi)$ is nontrivial).

Theorem 8.3. *Let $A = \{A_x\}_{x \in H}$ be a Hopf χ -coalgebra over an algebraically closed field \mathbb{k} such that the \mathbb{k} -algebra A_1 is semisimple and for all $x \in H$, the \mathbb{k} -algebra A_x is nonzero and finite dimensional. Then the category $\text{mod}_\chi(A)$ from Corollary 8.2 is χ -fusion.*

Proof. By [19, Lemma 5.1] applied to the Hopf H -coalgebra underlying A , we obtain that for all $x \in H$, the \mathbb{k} -algebra A_x is semisimple, and so the category of finite dimensional A_x -modules is semisimple. Note that all the irreducible A_x -modules are simple objects in the above sense because \mathbb{k} is an algebraically closed field, and there are at least one and only finitely many of them (up to isomorphism) since A_x is nonzero and finite dimensional. Also the unit object (which is \mathbb{k} with A_1 -action given by the counit) is an irreducible A_x -module. Then $\text{mod}_\chi(A)^1 = \text{mod}_H(A)$ is H -fusion.

It remains to verify Axiom (b). Let $e \in E$ and X be an object of $\text{mod}_\chi(A)$. Since the 1-subcategory $\text{mod}_\chi(A)^1$ is H -fusion, the object X is a 1-direct sum of a finite family $(i_\lambda)_{\lambda \in \Lambda}$ of homogeneous simple objects of $\text{mod}_\chi(A)^1$. It follows from the definition of $\text{mod}_\chi(A)$ that if i is a homogeneous simple object of $\text{mod}_\chi(A)^1$, then $(\phi_{\chi(e^{-1})|i|,e})^*(i)$ is a homogeneous simple object of $\text{mod}_\chi(A)^1$ which is e -isomorphic to i . Consequently, the object X is the e -direct sum of the finite family $((\phi_{\chi(e^{-1})|i_\lambda|,e})^*(i_\lambda))_{\lambda \in \Lambda}$ of homogeneous simple objects of $\text{mod}_\chi(A)^1$. \square

8.8. Representations of Hopf χ -algebras

Recall from Section 7.8 the notions of a χ -bialgebra and of a Hopf χ -algebra. Any χ -bialgebra $A = \bigoplus_{x \in H} A_x$ (over \mathbb{k}) yields an E -additive abelian χ -graded monoidal category $\text{Comod}_\chi(A)$. The construction of $\text{Comod}_\chi(A)$ is dual to that given in Section 8.2 and we only give here a brief description of it.

The objects of $\text{Comod}_\chi(A)$ are the (left) A -comodules, that is, the H -graded \mathbb{k} -module $M = \bigoplus_{x \in H} M_x$ such that each M_x is endowed with a structure of a (left) A_x -comodule. An A -comodule M is homogeneous of degree $x \in H$ if it is nonzero and is concentrated in degree x (meaning that $M = M_x$). The monoidal product of A -comodules is induced by the usual tensor product of H -graded \mathbb{k} -modules and the product $\mu = \{\mu_{x,y} : A_x \otimes A_y \rightarrow A_{xy}\}_{x,y \in H}$ of A .

For all $M, N \in \text{Comod}_\chi(A)$ and $e \in E$, the \mathbb{k} -module $\text{Hom}_{\text{Comod}_\chi(A)}^e(M, N)$ is the set of \mathbb{k} -linear homomorphisms $\alpha : M \rightarrow N$ such that for all $x \in H$,

- (a) $\alpha(M_x) \subset N_{\chi(e)x}$,
- (b) the map $(\phi_{x,e})_*(M_x) \rightarrow N_{\chi(e)x}$ induced by α is $A_{\chi(e)x}$ -colinear, where ϕ is the χ -action of A .

The composition is induced in the obvious way from the set-theoretical composition. The monoidal product of morphisms is their tensor product over \mathbb{k} . The monoidal constraints of $\text{Comod}_\chi(A)$ are induced (in the obvious way) from those of $\text{Mod}_{\mathbb{k}}$.

Similarly, the full subcategory $\text{comod}_\chi(A)$ of $\text{Comod}_\chi(A)$ consisting of the A -comodules whose underlying \mathbb{k} -module is projective of finite rank is an E -additive χ -graded monoidal category.

Dually to Theorem 8.1 and Corollary 8.2, if A is a Hopf χ -algebra, then the χ -category $\text{Comod}_\chi(A)$ is closed and the χ -category $\text{comod}_\chi(A)$ is rigid. Also, the dual $A^* = \{A_x^*\}_{x \in H}$ of a Hopf χ -algebra A of finite

type is a Hopf χ -coalgebra (see Section 7.8) and the χ -graded monoidal categories $\text{mod}_\chi(A^*)$ and $\text{comod}_\chi(A)$ are isomorphic. Conversely, the dual $A^* = \bigoplus_{x \in H} A_x^*$ of a Hopf χ -coalgebra $A = \{A_x\}_{x \in H}$ of finite type is a Hopf χ -algebra and the χ -graded monoidal categories $\text{comod}_\chi(A^*)$ and $\text{mod}_\chi(A)$ are isomorphic.

9. Hopf crossed module-modules and integrals

Throughout this section, $\chi: E \rightarrow H$ is a crossed module and $A = \{A_x\}_{x \in H}$ is a Hopf χ -coalgebra (over \mathbb{k}). We introduce Hopf χ -modules over A and prove a structure theorem for them. Next we use this theorem to prove the existence and uniqueness of χ -integrals for A .

9.1. Hopf χ -modules

A (left-left) Hopf χ -module over the Hopf χ -coalgebra A is a family $M = \{M_x\}_{x \in H}$ of \mathbb{k} -modules endowed with three families

$$\begin{aligned} r &= \{r_x: A_x \otimes M_x \rightarrow M_x\}_{x \in H}, \\ \rho &= \{\rho_{x,y}: M_{xy} \rightarrow A_x \otimes M_y\}_{x,y \in H}, \\ \psi &= \{\psi_{x,e}: M_x \rightarrow M_{\chi(e)x}\}_{(x,e) \in H \times E}, \end{aligned}$$

of \mathbb{k} -linear homomorphisms such that:

- (a) For all $x \in H$, the pair (M_x, r_x) is a (left) A_x -module.
- (b) The pair (M, ρ) is a (left) A -comodule, that is, for all $x, y, z \in H$,

$$(\Delta_{x,y} \otimes \text{id}_{M_z})\rho_{xy,z} = (\text{id}_{A_x} \otimes \rho_{y,z})\rho_{x,yz} \quad \text{and} \quad (\varepsilon \otimes \text{id}_{M_x})\rho_{1,x} = \text{id}_{M_x}.$$

- (c) The action r and coaction ρ intertwine as follows: for all $x, y \in H$,

$$\rho_{x,y}r_{xy} = (\mu_x \otimes r_y)(\text{id}_{A_x} \otimes \sigma_{A_y, A_x} \otimes \text{id}_{M_y})(\Delta_{x,y} \otimes \rho_{x,y}).$$

- (d) For all $x, y \in H$ and $e, f \in E$,

$$\begin{aligned} \psi_{x,1} &= \text{id}_{M_x}, & \psi_{e,x}r_x &= r_{\chi(e)x}(\phi_{x,e} \otimes \psi_{x,e}), \\ \psi_{x,fe} &= \psi_{\chi(e)x,f}\psi_{x,e}, & (\phi_{x,e} \otimes \psi_{y,f})\rho_{x,y} &= \rho_{\chi(e)x,\chi(f)y}\psi_{xy,ef}. \end{aligned}$$

Here $\mu = \{\mu_x\}_{x \in H}$, $\Delta = \{\Delta_{x,y}\}_{x,y \in H}$, ε , and $\phi = \{\phi_{x,e}\}_{(x,e) \in H \times E}$ are the product, coproduct, counit, and χ -action of A , respectively. Note that $\bigoplus_{x \in H} M_x$ is then an A -module in the sense of Sections 6.5 and 8.1.

9.2. Morphisms of Hopf χ -modules

Consider two Hopf χ -modules $M = (M, r, \rho, \psi)$ and $M' = (M', r', \rho', \psi')$ over A . A morphism of Hopf χ -modules from M to M' is a family $\theta = \{\theta_x: M_x \rightarrow M'_x\}_{x \in H}$ of \mathbb{k} -linear homomorphisms such that each θ_x is A_x -linear, θ is a morphism of A -comodules, and θ is a χ -equivariant:

$$r'_x(\text{id}_{A_x} \otimes \theta_x) = \theta_x r_x, \quad (\text{id}_{A_x} \otimes \theta_y)\rho_{x,y} = \rho'_{x,y}\theta_{xy}, \quad \theta_{\chi(e)x}\psi_{x,e} = \psi'_{x,e}\theta_x$$

for all $x, y \in H$ and $e \in E$.

Clearly the composition (componentwise) of two morphisms of Hopf χ -modules is a morphism of Hopf χ -modules.

9.3. Modules of coinvariants

The *module of coinvariants* of a Hopf χ -module $M = (M, r, \rho, \psi)$ over A is the \mathbb{k} -submodule $M^{\text{co}A}$ of $\prod_{x \in H} M_x$ consisting of the elements $m = (m_x)_{x \in H}$ such that for all $x, y \in H$ and $e \in E$,

$$\rho_{x,y}(m_{xy}) = 1_x \otimes m_y \quad \text{and} \quad \psi_{x,e}(m_x) = m_{\chi(e)x},$$

where 1_x is the unit element of A_x .

Any morphism of Hopf χ -modules $\theta: M \rightarrow M'$ induces a \mathbb{k} -linear homomorphism $\theta^{\text{co}A}: M^{\text{co}A} \rightarrow (M')^{\text{co}A}$ defined by

$$\theta^{\text{co}A}((m_x)_{x \in H}) = (\theta_x(m_x))_{x \in H}.$$

9.4. Trivial Hopf χ -modules

Let V be a \mathbb{k} -module. Then

$$(\{A_x \otimes V\}_{x \in H}, \{\mu_x \otimes \text{id}_V\}_{x \in H}, \{\Delta_{x,y} \otimes \text{id}_V\}_{x,y \in H}, \{\phi_{x,e} \otimes \text{id}_V\}_{(x,e) \in H \times E})$$

is a Hopf χ -module over A , denoted $A \otimes V$. Its module of coinvariants is

$$(A \otimes V)^{\text{co}A} = \{(1_x \otimes v)_{x \in H} \mid v \in V\}.$$

Note that the assignment $v \mapsto (1_x \otimes v)_{x \in H}$ is a \mathbb{k} -linear isomorphism $V \cong (A \otimes V)^{\text{co}A}$ and its inverse is computed by $(m_x)_{x \in H} \mapsto (\varepsilon \otimes \text{id}_V)(m_1)$.

Clearly, if $\alpha: V \rightarrow W$ is a \mathbb{k} -linear homomorphism, then $\{\text{id}_{A_x} \otimes \alpha\}_{x \in H}$ is a morphism of Hopf χ -modules from $A \otimes V$ to $A \otimes W$.

9.5. Structure of Hopf χ -modules

Let $\text{Mod}_{\mathbb{k}}$ be the category of \mathbb{k} -modules and \mathbb{k} -linear homomorphisms. Denote by ${}^A_A\mathcal{H}$ the category of Hopf χ -modules over A and their morphisms. The trivial Hopf χ -modules (see Section 9.4) and the modules of coinvariants (see Section 9.3) induce the following functors:

$$A \otimes ? : \text{Mod}_{\mathbb{k}} \rightarrow {}^A_A\mathcal{H} \quad \text{and} \quad ?^{\text{co}A} : {}^A_A\mathcal{H} \rightarrow \text{Mod}_{\mathbb{k}}.$$

Theorem 9.1. *The functors $A \otimes ?$ and $?^{\text{co}A}$ are equivalences and are quasi-inverse of each other. In particular, any Hopf χ -module M over A is isomorphic to the trivial Hopf χ -module $A \otimes M^{\text{co}A}$.*

Proof. It follows from the definitions that $?^{\text{co}A}$ is right adjoint to $A \otimes ?$ with unit $\eta_V: V \rightarrow (A \otimes V)^{\text{co}A}$ and counit $\epsilon_M = \{\epsilon_M^x: A_x \otimes M^{\text{co}A} \rightarrow M_x\}_{x \in H}$ given by

$$\eta_V(v) = (1_x \otimes v)_{x \in H} \quad \text{and} \quad \epsilon_M^x(a \otimes (m_y)_{y \in H}) = r_x(a \otimes m_x).$$

By Section 9.4, the unit η is an isomorphism. Let us prove that the counit ϵ is also an isomorphism. Let M be any Hopf χ -module over A . To prove that ϵ_M is an isomorphism of Hopf χ -modules from $A \otimes M^{\text{co}A}$ to M , we exhibit its inverse: defining $\pi: M_1 \rightarrow M^{\text{co}A}$ as

$$\pi(m) = (r_x(S_x \otimes \text{id}_{M_x})\rho_{x^{-1},x}(m))_{x \in H},$$

where $S = \{S_x\}_{x \in H}$ is the antipode of A , it follows from the definitions that

$$\nu_M = \{\nu_M^x = (\text{id}_{A_x} \otimes \pi)\rho_{x,1} : M_x \rightarrow A_x \otimes M^{\text{co}A}\}_{x \in H}$$

is the inverse of ϵ_M . \square

9.6. Integrals

A *left* (respectively, *right*) χ -integral for A is a family of \mathbb{k} -linear forms $\lambda = (\lambda_x : A_x \rightarrow \mathbb{k})_{x \in H}$ such that for all $x, y \in H$ and $e \in E$,

- $(\text{id}_{A_x} \otimes \lambda_y)\Delta_{x,y} = \eta_x \lambda_{xy}$ (respectively, $(\lambda_x \otimes \text{id}_{A_y})\Delta_{x,y} = \eta_y \lambda_{xy}$),
- $\lambda_{\chi(e)x} \phi_{x,e} = \lambda_x$.

Here $\{\eta_x : \mathbb{k} \rightarrow A_x\}_{x \in H}$, $\Delta = \{\Delta_{x,y}\}_{x,y \in H}$, and $\phi = \{\phi_{x,e}\}_{(x,e) \in H \times E}$ are the unit maps, coproduct, and χ -action of A .

We denote by \mathcal{I}_A^l (respectively, \mathcal{I}_A^r) the set of left (respectively, right) χ -integrals for A . The sets \mathcal{I}_A^l and \mathcal{I}_A^r are \mathbb{k} -modules (as submodules of the \mathbb{k} -module $\prod_{x \in H} A_x^*$). They are isomorphic: it follows from the properties of the antipode S of A (see Section 6.3 and Lemma 7.1) that the map

$$\mathcal{I}_A^l \rightarrow \mathcal{I}_A^r, \quad \lambda = (\lambda_x)_{x \in H} \mapsto \lambda^S = (\lambda_x^S = \lambda_{x^{-1}} S_{x^{-1}})_{x \in H}$$

is a \mathbb{k} -linear isomorphism.

Note that a χ -integral for A is in particular an H -integral (in the sense of [19]) for the Hopf H -coalgebra underlying A . (An H -integral for a Hopf H -coalgebra verifies only the first of the above two axioms of a χ -integral.) Consequently, by [19, Lemma 3.1], if a left or right χ -integral $\lambda = (\lambda_x)_{x \in H}$ for A is non-zero, then $\lambda_x \neq 0$ for all $x \in H$ such that $A_x \neq 0$, and in particular $\lambda_1 \neq 0$.

9.7. Existence and uniqueness of χ -integrals


It is known (see [15]) that the space of left (respectively, right) integrals for a finite dimensional Hopf algebra over a field is one dimensional. We generalize this result to Hopf χ -coalgebras:

Theorem 9.2. *Assume that \mathbb{k} is a field and that A is a Hopf χ -coalgebra (over \mathbb{k}) of finite type (that is, each A_x is finite dimensional). Then the spaces \mathcal{I}_A^l and \mathcal{I}_A^r are both one dimensional.*

Proof. For any $x, y \in H$ and $e \in E$, set $M_x = A_{x^{-1}}^* = \text{Hom}_{\mathbb{k}}(A_{x^{-1}}, \mathbb{k})$ and define the \mathbb{k} -linear homomorphisms

$$r_x : A_x \otimes M_x \rightarrow M_x, \quad \rho_{x,y} : M_{xy} \rightarrow A_x \otimes M_y, \quad \psi_{x,e} : M_x \rightarrow M_{\chi(e)x}$$

by



Here we use the convention of Section 2.7 for oriented arcs, a color $z \in H$ is an abbreviation for A_z (as in Section 6.4), and the dot represents the χ -action of A (see Section 7.1). It follows from the axioms of a Hopf χ -coalgebra and Lemma 7.1 that

$$M = (M = \{M_x\}_{x \in H}, r = \{r_x\}_{x \in H}, \rho = \{\rho_{x,y}\}_{x,y \in H}, \psi = \{\psi_{x,e}\}_{(x,e) \in H \times E})$$

is a Hopf χ -module over A . Then, by Theorem 9.1, M is isomorphic to the trivial Hopf χ -module $A \otimes M^{\text{co}A}$. Now it follows from the definitions of ρ, ψ , and of a right χ -integral that the map

$$\mathcal{I}_A^r \rightarrow M^{\text{co}A}, \quad \lambda = (\lambda_x)_{x \in H} \mapsto \lambda^{\text{co}} = (\lambda_{x^{-1}})_{x \in H}$$

is a \mathbb{k} -linear isomorphism. Thus M is isomorphic to the trivial Hopf χ -module $A \otimes \mathcal{I}_A^r$. In particular the \mathbb{k} -vector spaces M_1 and $A_1 \otimes \mathcal{I}_A^r$ are isomorphic and so

$$\dim(M_1) = \dim(A_1 \otimes \mathcal{I}_A^r) = \dim(A_1) \dim(\mathcal{I}_A^r).$$

Now $\dim(M_1) = \dim(A_1)$ (since $M_1 = A_1^*$ and A_1 is finite dimensional) and $\dim(A_1) \neq 0$ (since $A_1 \neq 0$ because $\varepsilon(1_1) = 1_{\mathbb{k}} \neq 0$). Consequently $\dim(\mathcal{I}_A^r) = 1$. Using that \mathcal{I}_A^l is isomorphic to \mathcal{I}_A^r (see Section 9.6), we get that $\dim(\mathcal{I}_A^l) = 1$. \square

9.8. The distinguished χ -grouplike element

As in Theorem 9.2, we assume in this subsection that \mathbb{k} is a field and that A is a Hopf χ -coalgebra (over \mathbb{k}) of finite type.

Corollary 9.3. *There exists a unique χ -grouplike element $g = (g_x)_{x \in H}$ of A such that $(\text{id}_{A_x} \otimes \lambda_y) \Delta_{x,y} = g_x \lambda_{xy}$ for any right χ -integral λ of A and all $x, y \in H$.*

Proof. First notice that the space \mathcal{I}_A^r of right χ -integrals for A coincide with the space of right H -integrals for the Hopf H -coalgebra underlying A because both are one dimensional (by Theorem 9.2 and [19, Theorem 3.6]) and any right χ -integral is a right H -integral. Then, by [19, Lemma 4.1], there is a unique H -grouplike element $g = (g_x)_{x \in H}$ of A satisfying the condition of the corollary. Now it follows from the properties of the χ -action ϕ of A that $(\phi_{\chi(e)x, e^{-1}}(g_{\chi(e)x}))_{x \in H}$ is also an H -grouplike element of A satisfying the condition of the lemma. Then the uniqueness of such an H -grouplike element implies that $\phi_{\chi(e)x, e^{-1}}(g_{\chi(e)x}) = g_x$, and so $g_{\chi(e)x} = \phi_{x,e}(g_x)$, for all $x \in H$ and $e \in E$. Hence g is a χ -grouplike element. \square

The χ -grouplike element $g = (g_x)_{x \in H}$ of Corollary 9.3 is called the *distinguished χ -grouplike element* of A . Note that g_1 is the (usual) distinguished grouplike element of the Hopf algebra A_1 . Also g is the distinguished H -grouplike element of the Hopf H -coalgebra underlying A (see [19, Section 4]).

10. Hopf crossed module-(co)algebras as Hopf algebras

Throughout this section, we let $\mathcal{S} = (\mathcal{S}, \otimes, \mathbb{1})$ be a symmetric monoidal category. We first define Hopf crossed module-(co)algebras in \mathcal{S} and then interpret them as Hopf algebras in some symmetric monoidal category associated with \mathcal{S} . As a corollary, when \mathcal{S} is the category of \mathbb{k} -modules, we obtain that the Hopf crossed module-(co)algebras over \mathbb{k} considered above are Hopf algebras in some symmetric monoidal category.

10.1. Hopf crossed module-(co)algebra in symmetric monoidal categories

Let $\chi: E \rightarrow H$ be a crossed module. The notion of a Hopf χ -(co)algebra in \mathcal{S} can be defined in the exact same way as in Sections 7.2 and 7.8 by replacing \mathbb{k} -modules with objects of \mathcal{S} and \mathbb{k} -linear homomorphisms with morphisms in \mathcal{S} .

Explicitly, a Hopf χ -coalgebra in \mathcal{S} is a family $A = \{A_x\}_{x \in H}$ of algebras in \mathcal{S} endowed with a coproduct, a counit, an antipode, and a χ -action which are respectively:

- a family $\Delta = \{\Delta_{x,y}: A_{xy} \rightarrow A_x \otimes A_y\}_{x,y \in H}$ of algebra morphisms in \mathcal{S} ,
- an algebra morphism $\varepsilon: A_1 \rightarrow \mathbb{1}$ in \mathcal{S} ,
- a family $S = \{S_x: A_{x^{-1}} \rightarrow A_x\}_{x \in H}$ of isomorphisms in \mathcal{S} ,
- a family $\phi = \{\phi_{x,e}: A_x \rightarrow A_{\chi(e)x}\}_{(x,e) \in H \times E}$ of algebra isomorphisms in \mathcal{S} .

These data should satisfy the following axioms: the coproduct must be coassociative and counital as in Section 6.1, the antipode must satisfy the axiom of Section 6.3, and the χ -action must satisfy the axioms (2)-(4) of Section 7.1.

Similarly, a Hopf χ -algebra in \mathcal{S} is a family $A = \{A_x\}_{x \in H}$ of coalgebras in \mathcal{S} endowed with a product, a unit, an antipode, and a χ -action which are respectively:

- a family $\mu = \{\mu_{x,y}: A_x \otimes A_y \rightarrow A_{xy}\}_{x,y \in H}$ of coalgebra morphisms in \mathcal{S} ,
- a coalgebra morphism $\eta: \mathbb{1} \rightarrow A_1$ in \mathcal{S} ,
- a family $S = \{S_x: A_x \rightarrow A_{x^{-1}}\}_{x \in H}$ of isomorphisms in \mathcal{S} ,
- a family $\phi = \{\phi_{x,e}: A_x \rightarrow A_{\chi(e)x}\}_{(x,e) \in H \times E}$ of coalgebra isomorphisms in \mathcal{S} ,

and which satisfy the axioms given in Section 7.8.

Note that these notions are dual to each other: Hopf χ -algebras in \mathcal{S} bijectively correspond to Hopf χ -coalgebras in the opposite category $\mathcal{S}^{op} = (\mathcal{S}^{op}, \otimes, \mathbb{1})$ by exchanging the product and the coproduct and by replacing the χ -action ϕ with $\phi^{-1} = \{\phi_{x,e}^{-1} = \phi_{\chi(e)x, e^{-1}}\}_{(x,e) \in H \times E}$.

For example, let $\text{Mod}_{\mathbb{k}}$ be the symmetric monoidal category of \mathbb{k} -modules and \mathbb{k} -linear homomorphisms. Then the Hopf χ -coalgebras over \mathbb{k} (as defined in Section 7.2) are exactly the Hopf χ -coalgebras in $\text{Mod}_{\mathbb{k}}$. Likewise, the Hopf χ -algebras over \mathbb{k} (as defined in Section 7.8) are exactly the Hopf χ -algebras in $\text{Mod}_{\mathbb{k}}$ and also correspond to the Hopf χ -coalgebras in $(\text{Mod}_{\mathbb{k}})^{op}$.

10.2. The category $\mathcal{E}(\mathcal{S})$

We associate to \mathcal{S} a symmetric monoidal category $\mathcal{E}(\mathcal{S})$ defined as follows.

The objects of $\mathcal{E}(\mathcal{S})$ are the pairs (\mathcal{C}, F) where \mathcal{C} is a small category and $F: \mathcal{C} \rightarrow \mathcal{S}$ is a functor. A morphism from an object (\mathcal{C}, F) to an object (\mathcal{D}, G) is a pair (Γ, γ) where $\Gamma: \mathcal{D} \rightarrow \mathcal{C}$ is a functor and $\gamma = \{\gamma_Y: F\Gamma(Y) \rightarrow G(Y)\}_{Y \in \mathcal{D}}$ is a natural transformation (from $F\Gamma$ to G). The composition of $(\Gamma, \gamma): (\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ with $(\Lambda, \lambda): (\mathcal{D}, G) \rightarrow (\mathcal{K}, H)$ is defined by

$$(\Lambda, \lambda) \circ (\Gamma, \gamma) = (\Gamma\Lambda, \lambda\gamma_\Lambda) \quad \text{where} \quad \lambda\gamma_\Lambda = \{\lambda_Z \circ \gamma_{\Lambda(Z)}\}_{Z \in \mathcal{K}}.$$

The identity of (\mathcal{C}, F) is $(1_{\mathcal{C}}, \text{id}_F)$ where $1_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ is the identity functor and $\text{id}_F = \{\text{id}_{F(X)}\}_{X \in \mathcal{C}}$. The monoidal product of two objects (\mathcal{C}, F) and (\mathcal{D}, G) is defined by

$$(\mathcal{C}, F) \boxtimes (\mathcal{D}, G) = (\mathcal{C} \times \mathcal{D}, F \otimes G = \otimes \circ (F \times G): \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{S}).$$

The monoidal product of a morphism $(\Gamma, \gamma): (\mathcal{C}, F) \rightarrow (\mathcal{C}', F')$ with a morphism $(\Lambda, \lambda): (\mathcal{D}, G) \rightarrow (\mathcal{D}', G')$ is defined by

$$(\Gamma, \gamma) \boxtimes (\Lambda, \lambda) = (\Gamma \times \Lambda, \gamma \otimes \lambda = \{\gamma_X \otimes \lambda_Y\}_{(X,Y) \in \mathcal{C}' \times \mathcal{D}'}).$$

This yields a functor $\boxtimes: \mathcal{E}(\mathcal{S}) \times \mathcal{E}(\mathcal{S}) \rightarrow \mathcal{E}(\mathcal{S})$. Set $I = (\mathbf{1}, \underline{\mathbb{1}})$, where $\mathbf{1}$ is the trivial category with a single object $*$ and a single morphism id_* , and the functor $\underline{\mathbb{1}}: \mathbf{1} \rightarrow \mathcal{S}$ is defined by $\underline{\mathbb{1}}(*) = \mathbb{1}$. Then $\mathcal{E}(\mathcal{S}) = (\mathcal{E}(\mathcal{S}), \boxtimes, I)$ is a monoidal category (with the obvious monoidal constraints) and is symmetric with symmetry induced (in the obvious way) from the symmetries of \mathbf{Cat} (see Section 2.3) and \mathcal{S} .

Note that the forgetful functor $\mathcal{U}: \mathcal{E}(\mathcal{S}) \rightarrow \mathbf{Cat}^{\text{op}}$, defined by $\mathcal{U}(\mathcal{C}, F) = \mathcal{C}$ and $\mathcal{U}(\Gamma, \gamma) = \Gamma$, is strict monoidal and symmetric.

10.3. Hopf algebras in $\mathcal{E}(\mathcal{S})$

In the next theorem, we characterize Hopf crossed module-coalgebras in \mathcal{S} as Hopf algebras in $\mathcal{E}(\mathcal{S})$.

Theorem 10.1. *Hopf algebras in $\mathcal{E}(\mathcal{S})$ are in bijective correspondence with pairs (χ, A) where χ is a crossed module and A is a Hopf χ -coalgebra in \mathcal{S} .*

Proof. Let (\mathcal{C}, F) be a Hopf algebra in the category $\mathcal{E}(\mathcal{S})$. Then $\mathcal{C} = \mathcal{U}(\mathcal{C}, F)$ is a Hopf algebra in \mathbf{Cat}^{op} . By Section 4.3 and since Hopf algebras in \mathbf{Cat}^{op} coincide with Hopf algebras in \mathbf{Cat} (by exchanging product and coproduct), there is a unique crossed module $\chi: E \rightarrow H$ such that $\mathcal{C} = (\mathcal{G}_\chi, \Delta_\chi, \varepsilon_\chi, m_\chi, u_\chi, S_\chi)$ as Hopf algebras in \mathbf{Cat}^{op} . The functor $F: \mathcal{G}_\chi \rightarrow \mathcal{S}$ gives rise to the family $A = \{A_x = F(x)\}_{x \in H}$ of objects in \mathcal{S} and to the family

$$\phi = \left\{ \phi_{x,e} = F(x \xrightarrow{e} \chi(e)x): A_x \rightarrow A_{\chi(e)x} \right\}_{(x,e) \in H \times E}$$

of morphisms in \mathcal{S} . The functoriality of F amounts to Axioms (2) and (3). The product of the Hopf algebra (\mathcal{C}, F) is (Δ_χ, μ) where $\mu = \{\mu_x: A_x \otimes A_x \rightarrow A_x\}_{x \in H}$ is a natural transformation from $(F \otimes F)\Delta_\chi$ to F . The unit of (\mathcal{C}, F) is (ε_χ, η) where $\eta = \{\eta_x: \mathbb{1} \rightarrow A_x\}_{x \in H}$ is a natural transformation from $\underline{\mathbb{1}}\varepsilon_\chi$ to F . The coproduct of (\mathcal{C}, F) is (m_χ, Δ) where $\Delta = \{\Delta_{x,y}: A_{xy} \rightarrow A_x \otimes_k A_y\}_{x,y \in H}$ is a natural transformation from Fm_χ to $F \otimes F$. The counit of (\mathcal{C}, F) is (u_χ, ε) where ε is a natural transformation from Fu_χ to $\underline{\mathbb{1}}$, that is, a morphism $\varepsilon: A_1 \rightarrow \mathbb{1}$ in \mathcal{S} . The antipode of (\mathcal{C}, F) is (S_χ, S) where $S = \{S_x: A_{x^{-1}} \rightarrow A_x\}_{x \in H}$ is a natural transformation from FS_χ to F . The associativity and unitality of the product of (\mathcal{C}, F) gives that each A_x is an algebra in \mathcal{S} with unit η_x . The naturality of the product and the unit of (\mathcal{C}, F) give that each $\phi_{x,e}$ is an algebra morphism. The coassociativity and counitality of the coproduct of (\mathcal{C}, F) give that Δ is coassociative and counital. The naturality of the coproduct of (\mathcal{C}, F) amounts to Axiom (4). The naturality of the counit is automatic. The fact that (\mathcal{C}, F) is a bialgebra in $\mathcal{E}(\mathcal{S})$ implies that each $\Delta_{x,y}$ and ε are algebra morphisms. The fact that (S_χ, S) is an invertible antipode for (\mathcal{C}, F) implies that each S_x is invertible and satisfies the axiom of Section 6.3. (Note that the naturality of the antipode of (\mathcal{C}, F) amounts to the property of Lemma 7.1 extended to Hopf χ -coalgebras in \mathcal{S} , and so can be deduced from the other axioms.) Then A is a Hopf χ -coalgebra in \mathcal{S} .

Conversely, any Hopf χ -coalgebra $A = \{A_x\}_{x \in H}$ in \mathcal{S} with χ -action ϕ gives rise to a Hopf algebra (\mathcal{G}_χ, F_A) in $\mathcal{E}(\mathcal{S})$, where the functor $F_A: \mathcal{G}_\chi \rightarrow \text{Mod}_k$ is defined by $F_A(x) = A_x$ and $F_A(x \xrightarrow{e} \chi(e)x) = \phi_{x,e}$, with (co)product, (co)unit, and antipode derived from those of A as above. \square

10.4. Hopf algebras in $\mathcal{F}(\mathcal{S})$

We associate to \mathcal{S} another symmetric monoidal category $\mathcal{F}(\mathcal{S})$ defined as follows.

Objects of $\mathcal{F}(\mathcal{S})$ are pairs (\mathcal{C}, F) where \mathcal{C} is a small category and $F: \mathcal{C} \rightarrow \mathcal{S}$ is a functor. A morphism from (\mathcal{C}, F) to (\mathcal{D}, G) is a pair (Γ, γ) where $\Gamma: \mathcal{C} \rightarrow \mathcal{D}$ is a functor and γ is a natural transformation from F to $G\Gamma$. The composition, monoidal product, and symmetry of $\mathcal{F}(\mathcal{S})$ are defined in a way similar to $\mathcal{E}(\mathcal{S})$. Explicitly, the composition of $(\Gamma, \gamma): (\mathcal{C}, F) \rightarrow (\mathcal{D}, G)$ with $(\Lambda, \lambda): (\mathcal{D}, G) \rightarrow (\mathcal{K}, H)$ is

$$(\Lambda, \lambda) \circ (\Gamma, \gamma) = (\Lambda\Gamma, \lambda_\Gamma\gamma) \quad \text{where} \quad \lambda_\Gamma\gamma = \{\lambda_{\Gamma(X)} \circ \gamma_X\}_{X \in \mathcal{C}}.$$

The identity of (\mathcal{C}, F) is $(1_{\mathcal{C}}, \text{id}_F)$. The monoidal product of two objects (\mathcal{C}, F) and (\mathcal{D}, G) is defined by

$$(\mathcal{C}, F) \boxtimes (\mathcal{D}, G) = (\mathcal{C} \times \mathcal{D}, F \otimes G = \otimes \circ (F \times G): \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{S}).$$

The monoidal product of a morphism $(\Gamma, \gamma): (\mathcal{C}, F) \rightarrow (\mathcal{C}', F')$ with a morphism $(\Lambda, \lambda): (\mathcal{D}, G) \rightarrow (\mathcal{D}', G')$ is defined by

$$(\Gamma, \gamma) \boxtimes (\Lambda, \lambda) = (\Gamma \times \Lambda, \gamma \otimes \lambda = \{\gamma_X \otimes \lambda_Y\}_{(X,Y) \in \mathcal{C} \times \mathcal{D}}).$$

The unit object of $\mathcal{F}(\mathcal{S})$ is $I = (\mathbf{1}, \underline{1})$, see Section 10.2. The monoidal constraints and symmetry of $\mathcal{F}(\mathcal{S})$ are induced (in the obvious way) from those of \mathbf{Cat} and \mathcal{S} . In particular, the forgetful functor $\mathcal{U}: \mathcal{E}(\mathcal{S}) \rightarrow \mathbf{Cat}$, defined by $\mathcal{U}(\mathcal{C}, F) = \mathcal{C}$ and $\mathcal{U}(\Gamma, \gamma) = \Gamma$, is strict monoidal and symmetric.

Theorem 10.2. *Hopf algebras in $\mathcal{F}(\mathcal{S})$ are in bijective correspondence with pairs (χ, A) where χ is a crossed module and A is a Hopf χ -algebra in \mathcal{S} .*

Proof. Recall that \mathcal{C}^{op} denotes the category opposite to a category \mathcal{C} . The opposite of a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is the functor $F^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$ acting as F on objects and morphisms. The opposite of a natural transformation α from a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ to a functor $G: \mathcal{C} \rightarrow \mathcal{D}$ is the natural transformation α^{op} from $F^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$ to $G^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$ defined by $\alpha_X^{\text{op}} = \alpha_X$ for all $X \in \text{Ob}(\mathcal{C}^{\text{op}}) = \text{Ob}(\mathcal{C})$.

Observe that the assignments $(\mathcal{C}, F) \mapsto (\mathcal{C}^{\text{op}}, F^{\text{op}})$ and $(\Gamma, \gamma) \mapsto (\Gamma^{\text{op}}, \gamma^{\text{op}})$ induce a symmetric strict monoidal isomorphism $\mathcal{F}(\mathcal{S}) \cong (\mathcal{E}(\mathcal{S}^{\text{op}}))^{\text{op}}$. Consequently, Hopf algebras in $\mathcal{F}(\mathcal{S})$ are in bijective correspondence with Hopf algebras in $(\mathcal{E}(\mathcal{S}^{\text{op}}))^{\text{op}}$, and so with Hopf algebras in $\mathcal{E}(\mathcal{S}^{\text{op}})$ (by exchanging product and coproduct), and so with pairs (χ, A) where χ is a crossed module and A is a Hopf χ -coalgebra in \mathcal{S}^{op} (by Theorem 10.1). We conclude using that Hopf χ -coalgebras in \mathcal{S}^{op} bijectively correspond to Hopf χ -algebras in the opposite category \mathcal{S} . \square

10.5. The case of Hopf crossed module-(co)algebras over \mathbb{k}

Applying Theorem 10.1 to the symmetric monoidal category $\text{Mod}_{\mathbb{k}}$ of \mathbb{k} -modules and \mathbb{k} -linear homomorphisms, we obtain the following characterization Hopf crossed module-coalgebras over \mathbb{k} as Hopf algebras:

Corollary 10.3. *Hopf algebras in $\mathcal{E}(\text{Mod}_{\mathbb{k}})$ are in bijective correspondence with pairs (χ, A) where χ is a crossed module and A is a Hopf χ -coalgebra over \mathbb{k} .*

Similarly, applying Theorem 10.2 to $\text{Mod}_{\mathbb{k}}$, we obtain the following characterization Hopf crossed module-algebras over \mathbb{k} as Hopf algebras:

Corollary 10.4. *Hopf algebras in $\mathcal{F}(\text{Mod}_{\mathbb{k}})$ are in bijective correspondence with pairs (χ, A) where χ is a crossed module and A is a Hopf χ -algebra over \mathbb{k} .*

Recall that if H is a group, then the trivial map $1 \rightarrow H$ is a crossed module and the notion of a Hopf $(1 \rightarrow H)$ -(co)algebra over \mathbb{k} corresponds to that of a Hopf H -(co)algebra over \mathbb{k} . Let $\mathcal{E}_d(\text{Mod}_{\mathbb{k}})$ and $\mathcal{F}_d(\text{Mod}_{\mathbb{k}})$

be the symmetric monoidal full subcategories of $\mathcal{E}(\text{Mod}_{\mathbb{k}})$ and $\mathcal{F}(\text{Mod}_{\mathbb{k}})$ consisting of the objects (\mathcal{C}, F) with \mathcal{C} a discrete small category (that is, a set). By restricting the correspondences of Corollaries 10.3 and 10.4 to these subcategories, we recover the following characterizations given in [4]: Hopf algebras in $\mathcal{E}_d(\text{Mod}_{\mathbb{k}})$ (respectively, in $\mathcal{F}_d(\text{Mod}_{\mathbb{k}})$) are in bijective correspondence with pairs (H, A) where H is a group and A is a Hopf H -coalgebra over \mathbb{k} (respectively, a Hopf H -algebra over \mathbb{k}).

CRedit authorship contribution statement

Kürşat Sözer: Writing – original draft, Writing – review & editing. **Alexis Virelizier:** Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported in part by the FNS-ANR grant OCHoTop (ANR-18-CE93-0002) and the Labex CEMPI (ANR-11-LABX-0007-01).

References

- [1] A. Bruguières, S. Lack, A. Virelizier, Hopf monads on monoidal categories, *Adv. Math.* 227 (2011) 745–800.
- [2] A. Bruguières, A. Virelizier, Hopf monads, *Adv. Math.* 215 (2007) 679–733.
- [3] R. Brown, C. Spencer, G-groupoids, crossed modules and the fundamental groupoid of a topological group, *Indag. Math.* 79 (1976) 296–302.
- [4] S. Caenepeel, M. De Lombaerde, A categorical approach to Turaev’s Hopf group-coalgebras, *Commun. Algebra* 34 (2006) 2631–2657.
- [5] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik, *Tensor Categories*, Mathematical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015.
- [6] M. Hennings, Invariants of links and 3-manifolds obtained from Hopf algebras, *J. Lond. Math. Soc.* 54 (1996) 594–624.
- [7] V. Jones, A polynomial invariant of knots via Von Neumann algebras, *Bull. Am. Math. Soc.* 12 (1985) 103–112.
- [8] A. Joyal, R. Street, The geometry of tensor calculus I, *Adv. Math.* 88 (1991) 55–112.
- [9] M. Kontsevich, Vassiliev’s knot invariants, *Adv. Sov. Math.* 16 (1993) 137–150.
- [10] G. Kuperberg, Noninvolutive Hopf algebras and 3-manifold invariants, *Duke Math. J.* 84 (1996) 83–129.
- [11] T. Le, J. Murakami, T. Ohtsuki, On a universal perturbative invariant of 3-manifolds, *Topology* 37 (1998) 539–574.
- [12] S. MacLane, *Categories for the Working Mathematician*, second edition, Springer-Verlag, New York, 1998.
- [13] N. Reshetikhin, V. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, *Invent. Math.* 103 (1991) 547–597.
- [14] K. Sözer, A. Virelizier, Monoidal categories graded by crossed modules and 3-dimensional HQFTs, *Adv. Math.* 428 (2023) 109155.
- [15] M.E. Sweedler, Integrals for Hopf algebras, *Ann. Math.* 89 (1969) 323–335.
- [16] V. Turaev, *Homotopy Quantum Field Theory*, EMS Tracts in Math., vol. 10, European Math. Soc. Publ. House, Zürich, 2010.
- [17] V. Turaev, A. Virelizier, *Monoidal Categories and Topological Field Theory*, Progress in Mathematics, vol. 322, Birkhäuser, Basel, 2017, xii+523 pp.
- [18] V. Turaev, O. Viro, State sum invariants of 3-manifolds and quantum 6j-symbols, *Topology* 31 (1992) 865–902.
- [19] A. Virelizier, Hopf group-coalgebras, *J. Pure Appl. Algebra* 171 (2002) 75–122.